



Letter to the Editor

Comment on “nutritional status and all-cause mortality in older adults with acute coronary syndrome”

Keywords:

Nutrition
Frailty
Mortality

Dear Editor,

We read the recent study with interest, published in *Clinical Nutrition*, by Tonet et al., in which they aimed to investigate the prognostic implication of nutritional status in older adults hospitalized for acute coronary syndrome [1]. In the study, they used the data of the populations of two different studies including “Frailty in Elderly Patients Receiving Cardiac Interventional Procedures” (FRASER) study and Impacto de la Fragilidad y otros Síndromes Geriátricos en el Manejo y Pronóstico Vital del Anciano con Síndrome Coronario Agudo sin Elevación de Segmento ST” (LONGEVO SCA) registry study [1]. They found that malnutrition, which was identified by Mini Nutritional Assessment-Short Form (MNA-SF), was an independent and strong risk factor for all-cause mortality. We think that some methodological issues should be considered while determining the relationship MNA scale or nutritional status and mortality in older adults [1].

Although malnutrition and frailty are two different major geriatric syndromes, it is known that these two syndromes, which can cause similar negative consequences, such as mortality, hospitalization, are in close relationship with each other [2]. Furthermore, in recent years, it has been shown that both MNA-long and MNA-SF can be a useful tool for frailty screening as well as malnutrition [3,4]. For example, Soysal P et al found that MNA-SF with a cut-off point of 11.0 had a sensitivity of 71.2% and a specificity of 92.8% for the detection of frail participants, and with a cut-off point of 13 had a sensitivity of 45.7% and a specificity of 78.3% for the detection of pre-frailty [4].

In the study by Tonet et al. used MNA-SF to identify nutritional status, and there were many frail and pre-frail participants in the two studies included (FRASER and LONGEVO SCA) [1]. This means that in the majority of patients with MNA-SF malnutrition and the risk of malnutrition, frailty and pre-frailty may occur simultaneously. Therefore, it is unclear whether the relationship

between nutritional status and mortality shown in this study is really due to malnutrition or frailty. In a similar study by Goldfarb et al., the one-year mortality risk for malnourished patients was approximately 3-fold higher than those with normal nutrition, but this rate was reduced to 1.08 when adjusted according to frailty [5]. Especially in the elderly, in order to establish a clear relationship between malnutrition and mortality, only non-frail individuals should be included in the study, or the potential impact of frailty on outcomes should be neutralized, as Goldfarb et al did.

In conclusion, the study contributes valuable data to medical literature, but the studies investigating nutritional status and clinical outcomes, such as mortality, should also evaluate frailty status, and make an adjustment for this, which in turn will provide a clearer picture to the readers.

Financial support

None declared.

Conflicts of interest

None declared.

Acknowledgements

There is no financial support and no conflict of interest.

References

- [1] Tonet E, Campo G, Maietti E, Formiga F, Martinez-Sellés M, et al. Nutritional status and all-cause mortality in older adults with acute coronary syndrome. *Clin Nutr* 2019 Jul 9. <https://doi.org/10.1016/j.clnu.2019.06.025> [in press].
- [2] Laur CV, McNicholl T, Valaitis R, Keller HH. Malnutrition or frailty? Overlap and evidence gaps in the diagnosis and treatment of frailty and malnutrition. *Appl Physiol Nutr Metabol* 2017 May;42(5):449–58.
- [3] Soysal P, Isik AT, Arik F, Kalan U, Eyvaz A, Veronese N. Validity of the mini-nutritional assessment scale for evaluating frailty status in older adults. *J Am Med Dir Assoc* 2019 Feb;20(2):183–7.
- [4] Soysal P, Veronese N, Arik F, Kalan U, Smith L, Isik AT. Mini Nutritional Assessment Scale-Short Form can be useful for frailty screening in older adults. *Clin Interv Aging* 2019 Apr 17;14:693–9.
- [5] Goldfarb M, Lauck S, Webb JG, Asgar AW, Perrault LP, Piazza N, et al. Malnutrition and mortality in frail and non-frail older adults undergoing aortic valve replacement. *Circulation* 2018;138:2202–11.

Pinar Soysal*

*Department of Geriatric Medicine, Bezmialem Vakif University,
Faculty of Medicine, Istanbul, Turkey*

Ahmet Turan Isik

*Unit for Aging Brain and Dementia, Department of Geriatric Medicine,
Dokuz Eylul University, Faculty of Medicine, Izmir, Turkey*

* Corresponding author. Adnan Menderes Bulvarı, Vatan Caddesi,
34093, Fatih, İstanbul, Turkey.

E-mail address: dr.pinarsoysal@hotmail.com (P. Soysal).

2 August 2019