

1-1-2015

Secondary metabolites from *Sambucus ebulus*

İREM ATAY

HASAN KIRMIZİBEKMEZ

AHMET CEYHAN GÖREN

ERDEM YEŞİLADA

Follow this and additional works at: <https://journals.tubitak.gov.tr/chem>

 Part of the [Chemistry Commons](#)

Recommended Citation

ATAY, İ, KIRMIZİBEKMEZ, H, GÖREN, A. C, & YEŞİLADA, E (2015). Secondary metabolites from *Sambucus ebulus*. *Turkish Journal of Chemistry* 39 (1): 34-41. <https://doi.org/10.3906/kim-1403-47>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Chemistry by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

Secondary metabolites from *Sambucus ebulus*

İrem ATAY^{1,*}, Hasan KIRMIZIBEKMEZ¹, Ahmet Ceyhan GÖREN²,
Erdem YEŞİLADA¹

¹Department of Pharmacognosy, Faculty of Pharmacy, Yeditepe University, Ataşehir, İstanbul, Turkey

²Chemistry Group Laboratories, TÜBİTAK UME, Gebze, Kocaeli, Turkey

Received: 17.03.2014 • Accepted: 19.06.2014 • Published Online: 23.01.2015 • Printed: 20.02.2015

Abstract: A new nonglycosidic iridoid, sambulin B (**1**), was isolated from the methanol extract of *Sambucus ebulus* L. leaves along with a recently reported new nonglycosidic iridoid, 10-*O*-acetylpatrinoside aglycone (sambulin A) (**2**); 2 flavonoids, isorhamnetin-3-*O*- β -D-glucopyranoside (**3**) and isorhamnetin-3-*O*-rutinoside (**4**); and a mixture of 2 flavonoids (**5**), quercetin-3-*O*- β -D-glucopyranoside and quercetin-3-*O*- β -D-galactopyranoside. Their structures were elucidated by 1-D and 2-D nuclear magnetic resonance spectroscopy (NMR) and mass spectrometry (MS) experiments.

Key words: *Sambucus ebulus*, nonglycosidic iridoids, sambulin B, flavonoids

1. Introduction

The genus *Sambucus* (Adoxaceae) is represented by 2 species in the flora of Turkey.¹ Among these, *Sambucus ebulus* L. is a perennial herb known locally as mürver, sultanotu, and şahmehlemi.^{1,2} In Anatolian folk medicine, *S. ebulus* is particularly used against inflammatory problems, i.e. rheumatic pain, edema, eczema, urticaria, burns, infectious wounds, and hemorrhoids as well as peptic ulcers.³ Anti-inflammatory, antinociceptive, wound-healing, cytotoxic, antiulcer, and anti-*Helicobacter pylori* effects of *S. ebulus* were reported in several previous studies.^{4–8} Simple phenols, flavonoids, anthocyanins, lignans, and iridoids have been shown to be the major secondary metabolites of the genus.^{9–13}

We herein present the isolation and structure elucidation of a new nonglycosidic iridoid named sambulin B (**1**), along with an iridoid, 10-*O*-acetylpatrinoside aglycone (sambulin A), and 4 known flavonoids, isorhamnetin-3-*O*- β -D-glucopyranoside (**3**), isorhamnetin-3-*O*-rutinoside (**4**), and a mixture of 2 flavonoids, hyperoside (quercetin-3-*O*- β -galactoside) and isoquercitrin (quercetin-3-*O*- β -glucoside) (**5**). The chemical structures of the compounds are presented in the Figure.

2. Results and discussion

The leaves of *S. ebulus* were extracted with MeOH. The crude extract was dispersed in 90% MeOH, and then submitted to partition with organic solvents in increasing polarity. Two nonglycosidic iridoids (**1** and **2**) were isolated from the *n*-hexane and CHCl₃ subextracts, while 2 flavonol glycosides and a mixture of 2 flavonoid glycosides were obtained from the EtOAc subextract.

*Correspondence: irematay@yahoo.com




Figure. Structures of the isolated compounds.

Compound **1** was obtained as a colorless oily substance from the *n*-hexane subextract. The molecular formula, $C_{19}H_{28}O_8$, was deduced from the pseudomolecular ion peaks at m/z 407 $[M+Na]^+$ and 791 $[2M+Na]^+$ in the ESI-MS and by inspecting ^{13}C NMR data. It exhibited UV maxima at 241, 274, and 282 nm. The IR spectrum suggested the presence of hydroxyl group (3678 cm^{-1}), olefinic C–H (3100 cm^{-1}), and ester carbonyl (1738 cm^{-1}) functionalities. The 1H NMR spectrum contained 1 olefinic (δ_H 6.34), 2 hydroxymethylene (δ_H 4.20 and 4.16, δ_H 4.09 and 4.00), 1 methylene (δ_H 2.15, 1.96), 1 oxymethyne (δ_H 5.30), 3 methyne (δ_H 3.01, 2.34, and 2.13), and 1 hemiacetal (δ_H 5.97) signals (Table 1). Moreover, the 1H NMR spectrum displayed 2 equivalent secondary methyl resonances at δ 0.97 (*d*, *J*: 6.4 Hz). In the COSY spectrum, 1 methyne resonance and 1 methylene (δ_H 2.26) signal conjugated to a carbonyl function were observed in a spin system. These data were indicative of the presence of an isovaleryl moiety. The signals at 22.3 (2C), 25.6, 42.9, and 171.7 in the ^{13}C NMR spectrum supported this assumption.¹⁴ Moreover, the presence of 2 acetoxy groups was evident from the signals at δ_H 2.05 (3H) and 2.04 (3H) and the corresponding carbon resonances at δ_C 21.0, 20.8, and 170.2 (2C). The ^{13}C NMR spectrum contained 19 signals; 5 of them were ascribed to an isovaleryl unit,

while 4 of them were arising from 2 acetyl groups. The remaining 10 carbon atoms were in accordance with a C₁₀-iridoid core. The esterification sites of the acyl units were established by the long-range correlations in the HMBC. Thus, cross peaks between carbonyl carbon of isovaleryl (δ_C 171.7) with H-1 (δ_H 5.97) and carbonyl carbon of one of the acetyl units with H₂-10 (δ_H 4.20, 4.16) revealed the locations of acyl units to be at C-1 and C-10, respectively. The NMR data of **1** were very close to those of 10-*O*-acetylpatrinoside, a recently published molecule from the same species by another research group, except for the presence of an additional acetyl unit in compound **1**. When the ¹H NMR spectrum of **1** was carefully inspected, H-7 signal of this appeared to be shifted to downfield about 1 ppm (δ_H 5.30) compared to 10-*O*-acetylpatrinoside aglycone (**2**).¹⁵ Moreover, the ¹³C NMR spectrum of the compound showed that the C-7 signal shifted downfield 2.5 ppm, while the C-6 and C-8 signals shifted upfield around 3 ppm compared to 10-*O*-acetylpatrinoside aglycone (**2**). Therefore, the position of the second acetyl group was determined to be C-7(OH). To determine the relative stereochemistry of the chiral centers in **1**, a ROESY experiment was performed. ROe cross-peaks were observed between H-1 α and H-8 α , and H-7 α and H-8 α , showing that these protons lie on the same side (α) of the molecule. In contrast, correlations were observed between H-3 and H-11 β , H-5 and H-9, H-5 and H-6 β , and H-5 and H-11 β , indicating that these protons were on the same side (β). Based on these findings the structure of compound **1** was elucidated as 7-*O*-acetyl derivative of 10-*O*-acetylpatrinoside aglycone (**2**). A literature survey revealed that compound **1** was a new nonglycosidic iridoid and was named sambulin B. On the other hand, for 10-*O*-acetylpatrinoside aglycone (**2**) we propose the trivial name sambulin A.

Table 1. ¹H (600 MHz, CDCl₃) and ¹³C-NMR (150 MHz, CDCl₃) data and HMBC correlations of sambulin B (1) and 10-*O*-acetylpatrinoside (sambulin A (2)).

	Sambulin B (1)		10- <i>O</i> -acetylpatrinoside aglycon (2)		HMBC (C→H)
Position	δ_C (ppm)	δ_H (ppm), <i>J</i> (Hz)	δ_C (ppm)	δ_H (ppm) <i>J</i> (Hz)	
1	91.1	5.97 <i>d</i> (4.7)	91.9	5.96 <i>d</i> (4.7)	H-3/H-8
3	138.4	6.34 <i>s</i>	138.0	6.39 <i>s</i>	H-1/H-5/H ₂ -11
4	116.9	-	118.0	-	H-3/H-5/H ₂ -6/H ₂ -11
5	32.0	3.01 <i>m</i>	32.5	3.07 <i>m</i>	H-1/H-3/H-7/H ₂ -11
6 α	37.2	2.15 <i>m</i>	39.6	2.19 <i>m</i>	H-5
6 β		1.96 <i>m</i>		1.70 <i>m</i>	
7	74.0	5.30 <i>m</i>	71.5	4.20 <i>m</i>	H-9/H ₂ -10
8	43.2	2.34 <i>m</i>	46.6	2.60 <i>m</i>	H ₂ -6
9	42.3	2.13 <i>m</i> †	41.1	2.16 <i>m</i> *	H ₂ -6/H-8/H ₂ -10
10a	62.2	4.20 <i>dd</i> (11.2, 6.9)	62.8	4.49 <i>dd</i> (11.1,6.9)	H-8
10b		4.16 <i>dd</i> (11.2, 6.9)		4.14 <i>dd</i> (11.1, 6.9)	
11a	62.6	4.09 <i>d</i> (12.3)	62.5	4.10 <i>d</i> (12.3)	H-3
11b		4.00 <i>d</i> (12.3)		4.02 <i>d</i> (12.3)	
7-COCH ₃	21.0	2.05 <i>s</i>			
7-COCH ₃	170.2	-			
10-COCH ₃	20.8	2.04 <i>s</i>	21.0	2.09 <i>s</i>	
10-COCH ₃	170.2	-	171.8	-	H ₂ -10/COCH ₃ (10)
1'	171.7	-	171.7	-	H-1/H ₂ -2'
2'	42.9	2.26 <i>m</i>	43.1	2.26 <i>m</i>	H ₃ -4'/H ₃ -5'
3'	25.6	2.30-2.22 <i>m</i> †	25.6	2.20-2.10 <i>m</i> †	H ₂ -2'/ H ₃ -4'/H ₃ -5'
4'	22.3	0.97 <i>d</i> (6.4)	22.4	0.98 <i>d</i> (6.4)	H ₂ -2'/H-3'
5'	22.3	0.97 <i>d</i> (6.4)	22.4	0.98 <i>d</i> (6.4)	H ₂ -2'/H-3'

*Unclear due to overlapping

In addition, 4 known flavonoids were isolated from the ethyl acetate subextract of *S. ebulus* methanol extract. The known compounds were identified by UV, IR, and 1-D and 2-D NMR techniques and by comparison with the previous published data. The structures of **3** and **4** were elucidated as isorhamnetin-3-*O*- β -D-glucopyranoside and isorhamnetin-3-*O*-rutinoside (Table 2). The NMR data of these compounds were found to be consistent with the published literature.¹⁶ Inspection of the ¹H NMR spectra of **5** revealed that it was a mixture of 2 flavonoid glycosides. A careful inspection of the spectra led to the identification of quercetin-3-*O*- β -D-glucopyranoside and quercetin-3-*O*- β -D-galactopyranoside. Thus **5** was identified as an inseparable mixture of hyperoside (quercetin-3-*O*- β -galactoside) and isoquercitrin (quercetin-3-*O*- β -glucoside). The ¹³C and ¹H NMR data of the hyperoside and isoquercitrin are given in Table 3; they were consistent with the previously published data.¹⁷

Table 2. ¹H (600 MHz, CD₃OD) and ¹³C NMR (150 MHz, CD₃OD) data of isorhamnetin-3-*O*- β -D-glucopyranoside (**3**) and isorhamnetin-3-*O*-rutinoside (**4**).

Position	Isorhamnetin-3- <i>O</i> - β -D-glucopyranoside (3)		Isorhamnetin-3- <i>O</i> -rutinoside (4)	
	δ_H , ppm (<i>J</i> , Hz)	δ_C , ppm	δ_H , ppm (<i>J</i> , Hz)	δ_C , ppm
2		157.2		159.2
3		130.9		132.9
4		178.0		179.9
5		160.5		164.0
6	6.21 <i>d</i> (2.1)	98.8	6.20 <i>d</i> (1.7)	100.7
7		165.1		163.5
8	6.41 <i>d</i> (2.1)	93.6	6.39 <i>d</i> (1.7)	95.6
9		157.0		159.0
10		104.1		104.9
1'		121.8		123.4
2'	7.91 <i>d</i> (2.1)	112.9	7.94 <i>d</i> (1.5)	116.4
3'		149.5		151.2
4'		147.0		148.9
5'	6.92 <i>d</i> (7.9)	114.7	6.90 <i>d</i> (8.5)	116.5
6'	7.58 <i>dd</i> (7.9, 2.1)	122.4	7.63 <i>dd</i> (8.5, 1.5)	124.5
1''	5.37 <i>d</i> (7.3)	100.2	5.22 <i>d</i> (7.6)	103.0
2''	3.24 <i>dd</i> (7.3, 5.2)	74.4	3.45-3.48 <i>m</i>	76.4
3''	3.46 <i>t</i> (7.3)	76.6	3.35-3.45 <i>m</i>	78.7
4''	3.30 <i>m</i>	69.9	3.35 <i>m</i>	72.1
5''	3.30 <i>m</i>	77.5	3.22 <i>m</i>	77.9
HA-6''	3.56 <i>dd</i> (12.0, 5.6)	61.2	3.55 <i>dd</i> (12.0, 5.1)	68.6
HB-6''	3.72 <i>dd</i> (12.0, 2.3)		3.72 <i>dd</i> (12.0, 2.5)	
OCH ₃	3.94 <i>s</i>	55.4	3.94 <i>s</i>	57.2
1'''			4.53 <i>d</i> (1.7)	102.9
2'''			3.20-3.80	72.4
3'''			3.20-3.80	72.8
4'''			3.20-3.80	74.4
5'''			3.20-3.80	70.3
6'''			1.1 <i>d</i> (6.5)	18.2

Several iridoids have been reported from *Sambucus* species.^{12,18,19} Recently 6 new 'Valeriana-type' iridoid glycosides were isolated from *S. ebulus* leaves. However, sambulin B is a new nonglycosidic ester iridoid, isolated from the leaves of *S. ebulus*. Since iridoids are accepted as significant chemotaxonomic markers, the occurrence of

such rare ‘Valeriana-type’ nonglycosidic iridoids might contribute to the chemotaxonomy of the genus *Sambucus* (formerly Caprifoliaceae) within its new family (Adoxaceae).

Table 3. ^1H (600 MHz, CD_3OD) and ^{13}C NMR (150 MHz, CD_3OD) data of quercetin-3-*O*- β -D-glucopyranoside (**5a**) and quercetin-3-*O*- β -D-galactopyranoside (**5b**).

Position	Quercetin-3- <i>O</i> - β -glucopyranoside (5a)		Quercetin-3- <i>O</i> - β -D-galactopyranoside (5b)	
	δ_{H} , ppm (<i>J</i> , Hz)	δ_{C} , ppm	δ_{H} , ppm (<i>J</i> , Hz)	δ_{C} , ppm
2		158.8		158.5
3		136.0		136.0
4		179.6		179.5
5		163.1		163.1
6	6.20 <i>d</i> (1.7)	99.9	6.20 <i>d</i> (2.1)	99.9
7		166.7		164.1
8	6.39 <i>d</i> (1.7)	94.8	6.39 <i>d</i> (2.1)	94.8
9		158.6		156.3
10		105.5		103.9
1'		123.1		123.1
2'	7.66 <i>d</i> (2.2)	116.1	7.84 <i>d</i> (2.3)	116.0
3'		145.9		145.9
4'		150.0		149.9
5'	6.84 <i>d</i> (8.5)	117.6	6.86 <i>d</i> (8.8)	117.8
6'	7.63 <i>dd</i> (8.5, 2.2)	123.2	7.59 <i>dd</i> (8.8, 2.3)	121.8
1''	5.25 <i>d</i> (7.6)	104.5	5.15 <i>d</i> (8.2)	102.3
2''	3.42 <i>dd</i> (7.6, 9.0)	75.8	3.53 <i>dd</i> (8.2, 9.3)	73.2
3''	3.48 <i>m</i>	78.4	3.22–3.34 <i>m</i>	75.1
4''	3.35 <i>m</i>	71.3	3.61 <i>d</i> (3.2)	70.1
5''	3.22 <i>m</i>	78.2	3.22–3.34 <i>m</i>	77.2
6''	3.58 <i>dd</i> (12.0, 5.5)	62.6	3.60*	61.9
	3.71 <i>dd</i> (12.0, 2.5)		3.71 <i>dd</i> (12, 2.5)	

*Unclear due to overlapping

3. Experimental

3.1. General

UV spectra were recorded on an HP Agilent 8453 (USA). IR spectra were recorded on a PerkinElmer-2000 FT-IR spectrometer (USA). NMR spectra were recorded on a Varian Mercury-Mx spectrometer (USA) at 600 MHz for ^1H NMR and at 150 MHz for ^{13}C NMR, with CDCl_3 and CD_3OD as solvents. Optical rotations were determined on an Opt. Act. Ltd. AA-5 polarimeter. Kieselgel 60 (0.063–0.200 mm; Merck, Darmstadt, Germany), Sephadex LH-20 (Lipophilic Sephadex, 25–100 μm , Sigma-Aldrich, USA), and Polyamide (Fluka, USA) were used for column chromatography (CC), and precoated Kieselgel 60 F₂₅₄ (Merck) plates were used for thin layer chromatography. For medium-pressure liquid chromatography (MPLC) a CombiFlash Companion (Teledyne Isco, USA) apparatus equipped with RediSep columns (C18, 130 and 43 g; Teledyne Isco, USA) was used.

3.2. Plant material

The leaves of *S. ebulus* were collected from Uludağ, Bursa (Turkey) in June 2009. The plants were identified by Prof Dr Erdem Yeşilada (Department of Pharmacognosy, Faculty of Pharmacy, Yeditepe University, İstanbul, Turkey). A voucher specimen (YEF 09017) was deposited at the Herbarium of Yeditepe University.

3.3. Extraction and isolation

The air dried and powdered leaves of *S. ebulus* (330 g) were extracted with methanol (MeOH) (2.3 l) over 24 h with intermittent stirring in a water bath (40 °C). The extract was filtered through filter paper and evaporated to dryness under reduced pressure to give crude MeOH extract (60.5 g, yield: 18.3%). The MeOH extract was then redissolved in 200 mL of MeOH in H₂O (10%) and extracted with *n*-hexane (9 × 200 mL). Combined *n*-hexane extract was evaporated under reduced pressure to yield *n*-hexane subextract (8 g, yield: 13.8%). Then MeOH was removed from the remaining extract and diluted with distilled H₂O and fractionated by successive solvent extractions with chloroform (CHCl₃) (4 × 200 mL), ethyl acetate (EtOAc) (4 × 200 mL), and *n*-butanol saturated with H₂O (*n*-BuOH) (4 × 100 mL). Each extract after solvent extraction was evaporated to dryness under reduced pressure to yield CHCl₃ subextract (22.5 g, yield: 39.2%), EtOAc subextract (2.5 g, yield: 4.3%), and *n*-BuOH subextract (9 g yield: 15.5%), and remaining water (11.5 g, yield: 19.8%).

3.3.1. Isolation of the components from EtOAc subextract

The EtOAc subextract (2 g) was further subjected to column chromatography (CC) over polyamide (25 g) and eluted with H₂O/MeOH mixtures in different ratios (0%–100%) to yield 4 fractions: SE_{EtOAc} I Fr. 1–4 (746 mg), SE_{EtOAc} I Fr. 5–10 (400 mg), SE_{EtOAc} I Fr. 12–14 (415 mg), and SE_{EtOAc} I Fr. 15–20 (227 mg). SE_{EtOAc} I Fr. 12–14 was subjected to MPLC over a RediSep Rf Reversed-phase C18 column (130 g) and eluted with mixtures of MeOH/H₂O (15%–100%) to yield 2 fractions: SE_{EtOAc} II Fr. 26–34 and SE_{EtOAc} II Fr. 35–49. Further CC of SE_{EtOAc} II Fr. 26–34 over SiO₂ (12 g) yielded a mixture (**5**) of hyperoside (quercetin-3-*O*-β-galactoside) and isoquercitrin (quercetin-3-*O*-β-glucoside) by eluting with mobile systems of MeOH/CHCl₃ (from 10% to 15%). The other fraction SE_{EtOAc} II Fr. 35–49 yielded isorhamnetin-3-*O*-β-D-glucopyranoside (**3**) by CC over SiO₂ (15 g) and eluting with mobile systems of MeOH/CHCl₃ (from 2% to 20%).

Furthermore, fractions of polyamide CC were further subjected to chromatographic separation by MPLC. SE_{EtOAc} I Fr. 15–20 was applied to MPLC over a RediSep flash column (12 g) and eluted with a MeOH/CHCl₃ solvent system (0%–80%). SE_{EtOAc} I Fr. 5–10 was applied to a MPLC RediSep 13 g reverse phase C18 column and eluted with MeOH/H₂O (5%–70%) to yield isorhamnetin-3-*O*-rutinoside (**4**).

3.3.2. Isolation of the components from CHCl₃ subextract

The CHCl₃ (10 g) subextract was applied to CC over SiO₂ (200 g) and eluted with first a solvent system composed of EtOAc/*n*-hexane in different rates (20%–100%) and then with MeOH/CHCl₃ mixtures (10%–50%) to give 4 fractions: SE_{CHCl3} I Fr. 2–7 (332 mg), SE_{CHCl3} I Fr. 8–13 (794 mg), SE_{CHCl3} I Fr. 14–16 (6.4 g), and SE_{CHCl3} I Fr. 17–23 (806 mg). SE_{CHCl3} I Fr. 14–16 was fractionated by CC over SiO₂ (150 g). Elution was started with CHCl₃ and then mixtures of MeOH/CHCl₃ (2.5%–60%) were used. Four subfractions were obtained: SE_{CHCl3} II Fr. 59–63 (1.289 g), SE_{CHCl3} II Fr. 71–72 (273 mg), SE_{CHCl3} II Fr. 73–74 (180 mg), and SE_{CHCl3} I Fr. 78–86 (615 mg). SE_{CHCl3} II Fr. 59–63 was applied to MPLC over a RediSep silica flash column (40 g) eluting with acetone/CHCl₃ (15%–90%) and 10-*O*-acetylpatrinoside aglycone (**2**) was obtained.

3.3.3. Isolation of the components from hexane subextract

The hexane subextract was applied to CC over Sephadex (16 g) and eluted with $\text{CHCl}_3/\text{MeOH}$ (50%) to yield 2 fractions: $\text{SE}_{\text{Hexane}}$ I Fr. 9–12 (38 mg) and $\text{SE}_{\text{Hexane}}$ I Fr. 13–15 (75 mg). The MPLC (RediSep Silica Flash Column (4 g)) of $\text{SE}_{\text{Hexane}}$ I Fr. 9–12 with $\text{EtOAc}/n\text{-hexane}$ (15%–80%) as eluent system yielded sambulin B (1).

3.4. Spectroscopic characteristics of the isolated compounds

Sambulin B (1): Colorless oily substance; $\text{C}_{19}\text{H}_{28}\text{O}_8$, $[\alpha]_D^{20} = +39.1$ ($c = 0.41$, CHCl_3); UV (CHCl_3): λ_{max} 241, 274, 282, IR (KBr) ν_{max} cm^{-1} : 3678, 3100, 1738, 1235, ESI-MS: $m/z = 407$ $[\text{M}+\text{Na}]^+$, 791 $[2\text{M}+\text{Na}]^+$, ^1H NMR (600 MHz, CDCl_3) and ^{13}C NMR (150 MHz, CDCl_3): Table 1.

10-O-acetylpatrinoside aglycone (sambulin A) (2): Colorless oily substance; $\text{C}_{17}\text{H}_{26}\text{O}_7$, $[\alpha]_D^{20} = -4.6$ ($c = 0.33$, CHCl_3); UV (CHCl_3): λ_{max} 241, 274, 282; IR (KBr) ν_{max} cm^{-1} : 3418, 1736, 1372, 1235; ESI-MS: $m/z = 343$ $[\text{M}+\text{H}]^+$, ^1H NMR (600 MHz, CDCl_3) and ^{13}C NMR (150 MHz, CDCl_3): Table 1.

Isorhamnetin-3-O- β -D-glucopyranoside (3): Yellow amorphous powder; $\text{C}_{22}\text{H}_{22}\text{O}_{12}$, UV (MeOH): λ_{max} 264, 354 nm, IR (KBr) ν_{max} cm^{-1} : 3409, 1653, 1607, HR-ESI-MS: $m/z = 501.1003$ $[\text{M}+\text{Na}]^+$, ^1H NMR (600 MHz, CD_3OD) and ^{13}C NMR (150 MHz, CD_3OD): Table 2.

Isorhamnetin-3-O-rutinoside (4): Yellow amorphous powder; $\text{C}_{28}\text{H}_{32}\text{O}_{16}$, UV (MeOH): λ_{max} 266, 354 nm, IR (KBr) ν_{max} cm^{-1} : 3452, 2934, 1665, 1608, HR-ESI-MS: $m/z = 647.1583$ $[\text{M}+\text{Na}]^+$, ^1H NMR (600 MHz, CD_3OD) and ^{13}C NMR (150 MHz, CD_3OD): Table 2.

Mixture of isoquercitrin (5a) and hyperoside (5b) (2/3) (5): Yellow amorphous powder; UV (MeOH): λ_{max} 266, 352 nm, IR (KBr) ν_{max} cm^{-1} : 3415, 1655, 1605, HR-ESI-MS: $m/z = 487.0847$ $[\text{M}+\text{Na}]^+$, ^1H NMR (600 MHz, CD_3OD) and ^{13}C NMR (150 MHz, CD_3OD): Table 3.

Acknowledgements

We greatly acknowledge the Scientific and Technological Research Council of Turkey (TÜBİTAK) for its support (SBAG 110S197).

References

1. Chamberline, D.; Phill, D.; Victoria, A. In *The Flora of Turkey and the East Aegean Islands Vol. 4*; Davis, P. H., Ed. Edinburgh Univ. Press: Edinburgh, UK, 1965–1988, pp. 541–543.
2. Yesilada, E.; Sezik, E.; Honda, G.; Takaishi, Y.; Takeda, Y.; Tanaka, T. *J. Ethnopharmacol.* **1999**, *64*, 195–210.
3. Sezik, E.; Zor, M.; Yesilada, E. *Int. J. Pharmacogn.* **1992**, *30*, 233–239.
4. Yesilada, E. *Chem. Nat. Comp.* **1997**, *5*, 539–540.
5. Suntar, I. P.; Akkol, E. K.; Yalcin, F. N.; Koca, U.; Keles, H.; Yesilada, E. *J. Ethnopharmacol.* **2010**, *129*, 106–114.
6. Ebrahimzadeh, M. A.; Mahmoudi, M.; Pourmorad, F.; Saeidnia, S.; Salimi, E. *J. Mazandaran Uni. Med. Sci.* **2006**, *16*, 35–42.
7. Shokrazadeh, M.; Mirzayi, M.; Saeedi, S. *Pharmacogn. Mag.* **2009**, *5*, 316–319.
8. Yesilada, E.; Gurbuz, I.; Toker, G. *J. Ethnopharmacol.* **2014**, *153*, 478–483.
9. Xiao, H. H.; Dai, Y.; Wan, H. Y.; Wong, M. S.; Yao, X. S. *British J. Nutr.* **2011**, *106*, 1802–1809.

10. Veberic, R.; Jakopic, J.; Stampar, F.; Schmitzer, V. *Food Chem.* **2009**, *114*, 511–515.
11. Yang, X. J.; Wong, M. S.; Wang, N. L.; Chan, S. C.; Yao, X. S. *J. Asian Nat. Prod. Res.* **2007**, *9*, 583–591.
12. Wang, Z. Y.; Han, H.; Yang, B. Y.; Xia, Y. G.; Kuang, H. X. *Molecules* **2011**, *16*, 3869–3874.
13. Sasaki, T.; Li, W.; Morimura, H.; Li, S.; Li, Q.; Asada, Y.; Koike, K. *Chem. Pharm. Bull.* **2011**, *59*, 1396–1399.
14. Pieri, V.; Schwaiger, S.; Ellmerer, E. P.; Stuppner, H. *J. Nat. Prod.* **2009**, *72*, 179–1803.
15. Tomassini, L.; Foddai, S.; Ventrone, A.; Nicoletti, M. *Nat. Prod. Res.* **2013**, *27*, 2012–2015.
16. Gutzeit, D.; Wray, V.; Winterhalter, P.; Jerz, G. *Chromatographia* **2007**, *1–2*, 1–7.
17. Olszewska, M. *Acta Poloniae Pharmaceutica* **2005**, *62*, 127–133.
18. Gross, G. A.; Sticher, O. *Helv. Chim. Acta* **1986**, *69*, 1113–1119.
19. Gross, G. A.; Sticher, O.; Anklin, C. *Helv. Chim. Acta* **1987**, *70*, 91–101.