

Curcumin Protects Against Ischemia/Reperfusion Injury in Rat Skeletal Muscle

Gulden Avci, M.D.,^{*,1} Huseyin Kadioglu, M.D.,[†] Ahmet O. Sehirli, M.D.,[‡] Suleyman Bozkurt, M.D.,[†] Oguz Guclu, M.D.,[§] Emrah Arslan, M.D.,^{*} and Sedit K. Muratli, M.D.[¶]

^{*}Department of Plastic and Reconstructive Surgery, Faculty of Medicine, Education and Research Hospital, Canakkale, Turkey; [†]Department of Surgery, Faculty of Medicine, Bezmialem University, Istanbul, Turkey; [‡]Department of Pharmacology, Faculty of Pharmacy, Marmara University, Istanbul, Turkey; [§]Department of Otorhinolaryngology, Faculty of Medicine, Canakkale Onsekiz Mart University, Canakkale, Turkey; and [¶]Department of Orthopedics and Traumatology, Faculty of Medicine, Canakkale Onsekiz Mart University, Canakkale, Turkey

Originally submitted May 3, 2011; accepted for publication August 10, 2011

Background. Curcumin has been shown to decrease ischemia-reperfusion (I/R) injury in kidney or brain tissues. In this study, the effects of curcumin were evaluated in skeletal muscle during I/R injury.

Materials and Methods. Hind limb ischemia was induced by clamping the common femoral artery and vein. After 4 h ischemia, the clamp of the femoral vessels of animals was taken off and the animal underwent 2 h reperfusion. We measured plasma concentrations of interleukin-1 β (IL-1 β) and tumor necrosis factor- α (TNF- α) using enzyme-linked immunosorbent assay (ELISA). The right gastrocnemius muscle was harvested and immediately stored at -30°C for the assessment of superoxide dismutase (SOD), catalase (CAT) activities, and measurement of glutathione (GSH), malondialdehyde (MDA), and protein oxidation (PO) levels. Curcumin (100 mg/kg), α -tocopherol, and normal saline (10 mL/kg) were administered intraperitoneally 1 h prior reperfusion.

Results. Plasma TNF- α or IL-1 β levels increased significantly in I/R group. The plasma levels of these proinflammatory cytokines were reduced in curcumin group. Muscle tissues of I/R groups revealed significantly higher antioxidant enzyme (superoxide dismutase, glutathione peroxidase, catalase) activities, and increased levels of malondialdehyde, nitric oxide, and protein carbonyl content compared with the SHAM group. Levels of these parameters in muscle revealed significant reductions in the I/R + curcumin

group compared with the I/R group. Curcumin has more potent antioxidant activity than vitamin E in the skeletal muscle I/R.

Conclusion. In this study, protective effects of curcumin against skeletal muscle ischemia-reperfusion injury have been revealed. We underscore the necessity of human studies with curcumin that would be hypothetically beneficial preventing skeletal muscle I/R injury. Crown Copyright © 2012 Published by Elsevier Inc. All rights reserved.

Key Words: ischemia-reperfusion injury; skeletal muscle; curcumin; reactive oxygen species.

INTRODUCTION

Ischemia and reperfusion (I/R) injury in skeletal muscle are unavoidable in many vascular and muscular traumas, diseases, and during a variety of time-consuming procedures or surgeries in the upper or lower extremities. Typical examples include limb amputation/replantation, transplantation, and free muscular-flap transfer. Ischemia reperfusion (I/R) injury is an influential factor that affects prognosis of cases subjected to reconstructive surgery, replantation surgery, and re-exploration. I/R injuries result in damage to endothelial and parenchymal cells, together with reactions of granulocytes and macrophages as well as humoral factors, including complement, coagulation factors, free radicals, and nitric oxide [1, 2]. I/R can activate macrophages and play a role in the release of cytokines which, in turn, influence leukocyte activation, transmigration, and target cell adhesion [3]. The enormous number of leukocytes that enter the extravascular space immediately after reperfusion release

¹ To whom correspondence and reprint requests should be addressed at Department of Plastic and Reconstructive Surgery, Faculty of Medicine, Canakkale Onsekiz Mart University Hospital for Education and Research, Kepez, Canakkale Turkey. E-mail: drguldenavci@hotmail.com.

large quantities of reactive oxygen species (ROS) capable of overwhelming cell antioxidant defenses [4, 5]. ROS, including hydroxyl radical (OH), superoxide anion radical (O²⁻), singlet oxygen (¹O²), hydrogen peroxide (H²O²) and nitric oxide (NO⁻) can cause cellular injury when they are generated excessively and hazardous to lipids, proteins, carbohydrates and nucleic acids [6–8]. Oxidative stress means an alteration in the delicate balance between free radicals and the scavenging capacity of antioxidant enzymes in favor of free radicals in the body systems [9]. ROS attack the polyunsaturated fatty acids in the membrane lipids, thereby lipid peroxidation results in loss of fluidity of the membranes and ruptures leading to release of cell [10]. I/R injury may extend beyond the ischemic area and damage the remote non-ischemic organs. For limiting or prevention of I/R injury, considerable effort has been made in developing new therapeutic strategies by the biotechnological and pharmaceutical industries. Many studies have been performed and various pharmacological and immunological agents have been used for this purpose [11–15].

The phytochemical curcumin (1,7-bis-[3-methoxy-4-hydroxyphenyl]-1,6-heptadiene-3,5-dione), a polyphenol derived from the root *Curcuma longa*, is a member of the ginger family. It has been used as a spice and food preservative for centuries. Curcumin has minimal side effects and has been administered safely in human subjects [16]. Curcumin's medicinal properties include antioxidant, antimicrobial, antitumor, antiproliferative, and anti-inflammatory properties [17–24]. Toda *et al.* showed that curcumin is a strong antioxidant compared with vitamins C and E [25].

In this study, we aimed to evaluate the antioxidant effects of curcumin against skeletal muscle I/R injury in a rat model. We analyzed the antioxidant enzymes (superoxide dismutase (SOD), catalase (CAT), glutathione (GSH), malondialdehyde (MDA), and protein oxidation (PO) levels in the muscle and, plasma levels of various inflammatory mediators, such as tumor necrosis factor- α (TNF- α), interleukin- β (IL-1 β).

MATERIALS AND METHODS

Forty Wistar rats, weighing between 200 and 250 g, were used. The animals were housed in temperature (21 \pm 2°C) and humidity (60% \pm 5%) controlled rooms in which a 12/12 h light/dark cycle was maintained with free access to food and tap water. All the protocols were approved by the institutional animal ethics committee of the Canakkale Onsekiz Mart University.

Experimental Design and Functional Studies

The animals were randomly divided into four groups: group 1 (sham, $n = 10$), the animals in this group were sham operated with the exposure of the femoral artery, but were not subjected to any ischemia reperfusion. group 2 (I/R+saline, $n = 10$), animals were ex-

posed to I/R, the rats in this group were subjected to 4 h of femoral pedicles occlusion followed by 2 h of reperfusion. Animals with I/R injury received saline (0.2 mL), with intraperitoneal injection 1 h before the reperfusion. Group 3 (curcumin+I/R, $n = 10$), animals were exposed to I/R, the rats in this group were subjected to 4 h of femoral pedicles occlusion followed by 2 h of reperfusion. Animals with I/R injury received curcumin (100 mg/kg) (Sigma, St. Louis, MO) with intraperitoneal injection 1 h before the reperfusion. Group 4 (vit E+I/R, $n = 10$), animals with I/R injury received α -tocopherol (Evigen-AKSU, Istanbul, Turkey) (10 mg/kg) with intraperitoneal injection 1 h before the reperfusion.

All surgical procedures were performed while the rats were under anesthesia with intraperitoneally administered 60 mg/kg ketamine and 10 mg/kg xylazine cocktail. Anesthesia was maintained by an additional half dose in every 90 min. Following induction of general anesthesia, the lower abdomen and groin were shaved, and sterile technique was employed in all surgical procedures. The animal was placed in a supine position. The right groin vessels were exposed via a transverse groin incision. The right femoral artery was isolated by clamping with an atraumatic microvascular clamp. After 4 h of ischemia, the microvascular clamp was removed. After removing the clamp, the lower extremity was inspected for restoration of blood flow. An ischemic insult was created in the right femoral artery for 4 h, followed by 2 h of reperfusion. Animals with I/R injury received either saline (0.2 mL), curcumin (100 mg/kg) (Sigma, St. Louis, MO) or α -tocopherol (Evigen-AKSU) (10 mg/kg) with intraperitoneal injection 1 h before the reperfusion. The dose of curcumin was based on previous studies [26]. A heat lamp was used to maintain the body temperature at 37 \pm 0.5°C during the experiment. At the end of the reperfusion period for biochemical assays blood was drawn from the inferior vena cava with heparinized syringes while still anesthetized. Blood samples were collected into polyethylene tubes and centrifuged (3500 \times g for 30 min at 4°C) to separate platelet-poor plasma. We measured TNF- α and IL 1 β in the plasma. Moreover, the right gastrocnemius muscle was harvested and immediately stored at -30°C for the assessment of SOD, CAT activities and measurement of GSH, MDA, and PO levels. Animals in the sham operation group underwent a surgical procedure similar to the other groups but the artery was not occluded.

Biochemical Assays

At the end of each experimental procedure, right gastrocnemius muscles were removed; the muscles were washed three times in cold isotonic saline (0.9% [vol/wt]). Then tissues were homogenized with cold Tris-HCl buffer (pH 7.4) to make a 10% homogenate. Glutathione measurements were taken using a modification of the Ellman procedure [27]. Briefly, after centrifugation at 2,000g for 10 min, 0.5 mL of supernatant was added to 2 mL of 0.3 mol/H₂PO₄-H₂O₂ solution. A 0.2 mL solution of dithiobisnitrobenzoate (0.4 mg/mL 1% sodium citrate) was added and the absorbance at 412 nm was measured immediately after mixing. Glutathione levels were calculated using an extinction coefficient of 1.36 \times 10⁵/M/cm. The results were expressed as nmol/100 mg protein in tissue samples. The total SOD activity was measured kinetically by a method described by Sun *et al.* [28]. The principle of the method is based on the inhibition of nitroblue tetrazolium reduction by the xanthine/xanthine oxidase system as a superoxide generator [28, 29]. One unit of SOD was defined as the enzyme amount causing 50% inhibition in the NBT reduction rate. SOD activity was expressed as units/mg of tissue protein.

Muscle tissue CAT (EC 1.11.1.6) activity was measurement according to Aebi's method [30]. The essential of the method was based on the determination of the rate constant k (s⁻¹, k) of the H²O² decomposition rate at 240 nm. Results were expressed as k (rate constant per gram protein; k g⁻¹ protein).

The MDA levels were assayed for products of lipid peroxidation by monitoring thiobarbituric acid reactive substance formation as described previously [31]. Lipid peroxidation is expressed in terms of

MDA equivalents using an extinction coefficient of 1.56×10^5 M/cm and the results are expressed as nmol MDA/g tissue.

The protein content of tissue samples was determined by the Lowry *et al.* assay [32]. Oxidized protein (PO) was quantified using the interaction between dinitrophenylhydrazine (DNP) and the carbonyls to yield a chromophore that absorbs strongly at 360 nm. All samples were diluted to 2–4 mg/mL of protein with wash buffer and treated with 1% streptomycin. The carbonyl content was calculated assuming a molar extinction coefficient of 22,000 [33]. The results were expressed as nmol/ μ g of tissue protein.

Serum levels of TNF- α and IL-1 β were quantified using enzyme-linked immunosorbent assay (ELISA) kits specific for the previously mentioned rat cytokines according to the manufacturer's instructions and guidelines (Biosource Europe SA, Nivelles, Belgium). The total AOC in plasma was measured using colorimetric test system (ImAnOx, catalog no. KC5200; Immunodiagnostic AG, Bensheim, Germany), according to the instructions provided by the manufacturer. The 8-hydroxy-2 α -deoxyguanosine (8-OHdG) content in the extracted DNA solution was determined by ELISA method (highly sensitive 8-OHdG ELISA kit; Japan Institute for the Control of Aging, Shizuoka, Japan). These particular assay kits were selected because of their high degree of sensitivity, specificity, inter- and intra-assay precision, and small amount of plasma sample required to conduct the assay. The concentration of TNF- α and IL 1 β in plasma were expressed as pg/mL.

Statistical Analysis

Statistical analysis was carried out using GraphPad Prism 3.0 (GraphPad Software, San Diego, CA). All data were expressed as mean \pm SEM. Groups of data were compared with an analysis of variance (ANOVA) followed by Tukey's multiple comparison tests. *P* values <0.05 were regarded as significant.

RESULTS

Results of oxidative stress markers in each group were shown in Tables 1 and 2. The GSH levels of the muscle tissue were as follows: sham group, 2.11 ± 0.18 ; control group, 0.94 ± 0.11 ; and curcumin treated group, 2.03 ± 0.12 ; and α -tocopherol treated group 1.83 ± 0.13 . The muscle tissue GSH levels in the control group were significantly lower than the sham and curcumin-treated group and α -tocopherol treated group ($P < 0.001$ and $P < 0.01$, respectively). The difference between the sham and curcumin treated group and α -tocopherol treated group was significant ($P < 0.05$; see Table 1). The difference be-

tween the curcumin and α -tocopherol treated groups was also significant (Fig. 1). Ischemia-reperfusion injury to the muscle significantly decreased SOD activity muscle tissues compared with the sham group. SOD levels were significantly improved by curcumin-treated and α -tocopherol treated groups and there were no significant differences in the curcumin-treated and α -tocopherol treated groups (Fig. 2).

CAT activity in muscle tissue decreased markedly after reperfusion compared with the sham group. A significant decrease in the activity of CAT was observed in I/R group compared with curcumin treated group and α -tocopherol treated group. The difference between the curcumin and α -tocopherol treated groups was also significant (Fig. 3).

The MDA levels of the muscle tissue were as follows: sham group, 18.30 ± 2.59 ; control group, 44.67 ± 5.81 ; curcumin-treated group, 19.39 ± 3.09 ; and α -tocopherol treated group 23.84 ± 4.83 . The tissue MDA levels in the control group were significantly higher than the sham and curcumin-treated and α -tocopherol treated groups. The difference between the sham and the curcumin and α -tocopherol treated groups was also significant ($P < 0.001$). The difference between the curcumin and α -tocopherol treated groups was also significant (Fig. 4). Carbonyl concentrations were 1.53 ± 0.17 nmol/mg proteins in sham group and 4.34 ± 0.51 nmol/mg protein in control group. In the curcumin treated group it was increased to 1.59 ± 0.16 nmol/mg protein ($P < 0.001$) and in α tocopherol treated rats it was increased to 2.21 ± 0.27 nmol/mg protein (Fig. 5).

The plasma levels of proinflammatory cytokines (TNF- α and IL-1 β) in the control (ischemia reperfusion injury in the skeletal muscle) group were significantly higher ($P < 0.001$) than that of the control group, whereas the treatment of curcumin and α -tocopherol abolished these elevations significantly ($P < 0.01$). The differences of the plasma levels of the TNF- α and IL-1 β between the curcumin and α -tocopherol treated groups was also significant (Fig. 6).

TABLE 1

Comparison of Effects of Ischemia/Reperfusion (I/R) on the Muscle GSH, SOD, CAT, MDA, PO Levels with the Curcumin-+IR, vit e+ I/R, Control and Sham Groups

	Sham	I/R-SF control	I/R-curcumin	I/R- α -tocopherol
GSH (μ mol/g)	2.11 ± 0.18	$0.94 \pm 0.11^{***}$	$2.03 \pm 0.12^{\dagger\dagger\dagger}$	$1.83 \pm 0.13^{\dagger\dagger}$
SOD (U/mgprotein)	0.95 ± 0.11	$0.48 \pm 0.06^{**}$	$0.89 \pm 0.10^{\dagger}$	$0.88 \pm 0.05^{\dagger}$
Catalase (U/mg protein)	61.01 ± 6.35	$26.57 \pm 5.26^{**}$	$57.70 \pm 5.38^{\dagger\dagger}$	$52.25 \pm 6.44^{\dagger}$
MDA (nmol/g)	18.30 ± 2.59	$44.67 \pm 5.81^{**}$	$19.39 \pm 3.09^{\dagger\dagger}$	$23.84 \pm 4.83^{\dagger}$
PO (nmol carbonyl/mg)	1.53 ± 0.17	$4.34 \pm 0.51^{***}$	$1.59 \pm 0.16^{\dagger\dagger\dagger}$	$2.21 \pm 0.27^{\dagger\dagger\dagger}$

* $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ compared with the sham group.

$^{\dagger}P < 0.05$, $^{\dagger\dagger}P < 0.01$, $^{\dagger\dagger\dagger}P < 0.001$ compared with the I/R group.

TABLE 2

Comparison of Effects of Ischemia/Reperfusion (I/R) on the Muscle TNF- α and IL-1 β levels with the Curcumin-+IR, vit e+ I/R, Control and Sham Groups

	Control	I/R	I/R-curcumin	I/R- α -tocopherol
TNF- α (pg/mL)	7.63 \pm 1.14	33.98 \pm 3.65***	11.15 \pm 2.29 ^{†††}	16.17 \pm 2.97 ^{†††}
IL-1 β (pg/mL)	10.13 \pm 1.46	32.69 \pm 2.94***	12.89 \pm 1.98 ^{†††}	19.17 \pm 2.90 ^{†††}

*** $P < 0.001$ compared with the sham group.

^{†††} $P < 0.01$, ^{††††} $P < 0.001$ compared with the I/R group.

DISCUSSION

Restoration of blood flow is critical for reversing ischemia; paradoxically the insult to the tissues worsens [34]. After ischemia, tissues were shown to contain an accumulation of xanthine oxidase [35]. This enzyme uses molecular oxygen, which is available as the final electron acceptor. This reaction produces singlet oxygen molecules, which are extremely unstable. Using these molecules, secondary chemical reactions produce superoxide, hydrogen peroxide, and hydroxyl. ROS, particularly the hydroxyl radicals, interact with phospholipids, proteins, and nucleic acids leading to lipid peroxidation, which results in a loss of membrane integrity [34]. The post-ischemic endothelium is the main source of the superoxide radical. Indirectly, superoxide may be responsible for the production of the damaging hydroxyl radical. The tissue damage caused by the production of ROS can trigger several defense mechanisms. The first-line defense mechanism includes antioxidants such as superoxide dismutase (SOD), catalase (CAT), and glutathione peroxidase (GSH). These enzymes catalyze the conversion of ROS into less reactive species [34, 36]. Under normal conditions, the damaging effects of superoxide are prevented by SOD,

which converts superoxide into hydrogen peroxide [35, 37]. However, during reperfusion of ischaemic tissues, these natural defenses may be overcome and hydrogen peroxide is converted into a hydroxyl radical, which in turn is capable of damaging a wide variety of biological molecules, including amino acids, membrane transport proteins, and nucleic acids [37, 38]. GSH is an important antioxidant, and the increased concentration of GSH in cells may be useful in the prevention of oxidative damage of endothelial cells [36]. GSH is part of the defensive strategy that cells and tissues use to combat oxidation. In the literature, there are several studies demonstrating that I/R in the tissue is associated with lipid peroxidation, which is an autocatalytic mechanism leading to oxidative destruction of cellular membranes [37]. Catalase, an oxidoreductase that catalyzes the conversion of hydrogen peroxide to water and oxygen, also can protect cells from damage induced by ischemia/reperfusion through scavenging ROS. GSH and SOD are the most important endogenous antioxidant defense mechanisms against oxidative stress [39, 40].

The most important damaging effect of free radicals on tissues is lipid peroxidation. The cell membrane is composed of fatty acids and phospholipids. Oxygen free radicals cause cellular injury by inducing lipid peroxidation, which results in functional and structural




FIG. 1. The effects of ischemia/reperfusion (I/R) and the treatment with curcumin, the treatment with α tocopherol on the GSH levels. *** $P < 0.001$ compared with the control group. ^{††} $P < 0.01$ compared with the IR group.




FIG. 2. The effects of ischemia/reperfusion (I/R) and the treatment with curcumin, the treatment with α tocopherol on the SOD levels. ⁺ $P < 0.05$ compared with the control group. ** $P < 0.01$ compared with the IR group.




FIG. 3. The effects of ischemia/reperfusion (I/R) and the treatment with curcumin, the treatment with α tocopheral on the CAT levels. $^+P < 0.05$, $^{**}P < 0.001$ compared with the control group. $^{++}P < 0.01$ compared with the IR group.

cell alterations [41]. Furthermore, ROS cause oxidation of protein structures and the protein damage can be revealed by the measurement of tissues protein carbonyl content. Malondialdehyde, a product of lipid peroxidation, is generated as a result of active oxygen radicals, which destroy unsaturated fatty acids in cell membranes [42].

ROS during ischemia and reperfusion to the skeletal muscle have a number of features, which include impaired muscle contractions, muscle necrosis, endothelial cell swelling, release of cellular proteins, and increased microvascular permeability to proteins [43, 44]. Many studies have suggested the beneficial effects of nutrients or other agents such as vitamin E [45], vitamin C [46], green tea extract [45], melatonin [47, 48], carbenoxolone [49], and green propolis [50, 51] in reducing or preventing cerebral or muscle injury during ischemia-reperfusion.




FIG. 4. The effects of ischemia/reperfusion (I/R) and the treatment with curcumin, the treatment with α tocopheral on the MDA levels. $^+P < 0.05$, $^{++}P < 0.001$ compared with the control group. $^{**}P < 0.01$ compared with the IR group.




FIG. 5. The effects of ischemia/reperfusion (I/R) and the treatment with curcumin, the treatment with α tocopheral on the PO levels. $^{+++}P < 0.001$ compared with the control group. $^{***}P < 0.01$ compared with the IR group.

In this study, we employed a previously studied model of femoral vessels atraumatic clamping-induced skeletal muscle ischemia-reperfusion injury. Clamping-induced skeletal muscle ischemia-reperfusion injury is associated with free radical production and proinflammatory responses. Treatment with curcumin improves kidney,




FIG. 6. The effects of ischemia/reperfusion (I/R) and the treatment with curcumin, the treatment with α tocopheral on the TNF- α and IL-1 β levels. $^{+++}P < 0.001$ compared with the control group. $^{***}P < 0.01$ compared with the IR group.

myocardium, pancreas, liver, lung mesentery ischemia-reperfusion injury. The aim of this study was to assess the hypothesis that curcumin attenuates skeletal muscle ischemia-reperfusion injury. Moreover, in the present study, we compared the effects of curcumin with a potent antioxidant, i.e., vitamin E. Curcumin is an effective scavenger for reactive oxygen radicals and reactive nitrogen species *in vitro* [52]. However, it has not been clear whether curcumin acts directly as an antioxidant *in vivo*. Curcumin has been shown to decrease the number of oxidative DNA adducts in malignant colorectal tissue [53]. In addition to direct antioxidant activity, curcumin may function indirectly as an antioxidant by inhibiting the activity of inflammatory enzymes or by enhancing the synthesis of glutathione, an important intracellular antioxidant. Chan *et al.* [18] reported that *in vivo* curcumin treatment produced 50%–70% reductions in iNOS mRNA expression in the liver of lipopolysaccharide (LPS)-injected mice. It has been reported that low concentrations of curcumin could inhibit NO production in activated macrophages, demonstrating that it significantly reduces the mRNA levels and 130-kDa inducible NOS protein expressed in activated macrophages [54]. Additionally, Shen *et al.* [54] reported that curcumin pretreatment protected the liver from warm I/R injury through multiple pathways, regulating expression and activity of multiple bioactive molecules and inhibiting cell apoptosis and neutrophil infiltration. Overexpression of antioxidant enzymes after curcumin pretreatment may play an essential role to protect the liver against warm I/R injury [53, 54]. The protective effects of curcumin on ischemia-reperfusion induced lung inflammation, capillary barrier dysfunction, tissue edema, and injury, in a similar manner to the steroid dexamethasone [55]. Curcumin reduces ischemia-reperfusion induced acute lung injury, probably through improvement of oxidative stress and nuclear factor- κ B-mediated expression of inflammatory cytokines [53]. Curcumin, which has antioxidant and anti-inflammatory effects, significantly attenuated the inflammatory responses and airway hyper-reactivity induced by pancreatic I/R [56].

The administration of vitamin E prevents the muscles from oxidative stress, endothelial damage, intramuscular edema, and major muscle fiber damage [57, 58]. It has been demonstrated that vitamin E had a protective effect on oxidative muscle damage after a period of I/R in human. It was observed that MDA level and neutrophil infiltration in muscle fibers decreased in vitamin E-treated patients after a period of I/R [58]. Our results revealed that I/R injury led increases in MDA and PO levels in the skeletal muscle tissue. In our study, we showed that administration of curcumin significantly decreased tissue protein carbonyl levels as a marker of protein oxidation activities

(PO) in skeletal muscle compared with I/R group. Curcumin reversed the increase of MDA levels to a considerable extent, thereby confirming its antioxidant role in I/R. Reactive oxygen species attack the polyunsaturated fatty acids in the membrane lipids and result in peroxidation, which may lead to disorganization of cell structure and function. This process results in an excess of free radicals, which can react with cellular lipids, proteins, and nucleic acids, leading to local injury and eventually organ dysfunction. It has been suggested that the protective effects of curcumin depend mainly on its antioxidant properties.

In this study, we also demonstrated the effect of curcumin on tissue GSH levels and SOD, CAT activity in a hind limb model of I/R injury. We found decreased SOD, CAT and GSH activities in I/R group compared to sham group. A significant decrease in the activity of SOD, CAT, and GSH was observed in I/R group compared with curcumin treatment group. Furthermore, plasma levels of the proinflammatory cytokines TNF- α and IL-1 β were increased in I/R group compared with Sham group. Plasma levels of the proinflammatory cytokines TNF- α and IL-1 β were reduced in curcumin group. Moreover, in the present study, we compared the effects of curcumin with one of the potent antioxidants; herewith curcumin has more potent antioxidant activity than vitamin E in the skeletal muscle I/R.

In conclusion, this study demonstrated that administration of curcumin might protect skeletal muscle from reperfusion injury to the extent that vitamin E did. Our data also revealed that this protective effect of curcumin is probably ascribed to its free radical scavenging activity. We underscore the necessity of human studies with curcumin that would be hypothetically beneficial in preventing skeletal muscle I/R injury, particularly during surgical interventions.

REFERENCES

1. Land WG. The role of postischemic reperfusion injury and other nonantigen-dependent inflammatory pathways in transplantation. *Transplantation* 2005;79:505.
2. Behrend M. Immune-adhesion molecules in the prevention of allograft rejection and reperfusion injury. *Expert Opin Investig Drugs* 2000;9:789.
3. Le Moine O, Louis H, Stordeur P, et al. Role of reactive oxygen intermediates in interleukin 10 release after cold liver ischemia and reperfusion in mice. *Gastroenterology* 1997;113:1701.
4. Jaeschke H, Farhood A. Neutrophil and Kupffer cell-induced oxidant stress and ischemia-reperfusion injury in rat liver. *Am J Physiol* 1991;260:G355.
5. Hsieh YH, Huang SS, Wei FC, et al. Resveratrol attenuates ischemia-reperfusion induced leukocyte-endothelial cell adhesive interactions and prolongs allograft survival across the MHC barrier. *Circ J* 2007;71:423.
6. Clanton TL, Zuo L, Klawitter P. Oxidants and skeletal muscle function: Physiologic and pathophysiologic implications. *Proc Soc Exp Biol Med* 1999;222:253.

7. Cheeseman KH. Mechanisms and effects of lipid peroxidation. *Mol Aspects Med* 1993;14:191.
8. Marx G, Chevion M. Site-specific modification of albumin by free radicals. Reaction with copper (II) and ascorbate. *Biochem J* 1986;236:397.
9. Frei B. Reactive oxygen species and antioxidant vitamins: Mechanisms of action. *Am J Med* 1994;26:5S.
10. Ozyurt H, Ozyurt B, Koca K, et al. Caffeic acid phenethyl ester (CAPE) protects rat skeletal muscle against ischemia-reperfusion-induced oxidative stress. *Vascul Pharmacol* 2007; 47:108.
11. Portakal O, Inal-Erden M. Effects of pentoxifylline and coenzyme Q10 in hepatic ischemia/reperfusion injury. *Clin Biochem* 1999;32:461.
12. Huk I, Brovkovich V, Nanobashvili J, et al. Prostaglandin E1 reduces ischemia/reperfusion injury by normalizing nitric oxide and superoxide release. *Shock* 2000;14:234.
13. Calapai G, Marciano MC, Corica F, et al. Erythropoietin protects against brain ischemic injury by inhibition of nitric oxide formation. *Eur J Pharmacol* 2000;401:349.
14. Konukoglu D, Taşci I, Cetinkale O. Effects of cyclosporin A and ibuprofen on liver ischemia-reperfusion injury in the rat. *Clin Chim Acta* 1998;275:1.
15. Gelgor L, Butkow N, Mitchell D. Effects of systemic non-steroidal anti-inflammatory drugs on nociception during tail ischemia and on reperfusion hyperalgesia in rats. *Br J Pharmacol* 1992;105:412.
16. Chainani-Wu N. Safety and anti-inflammatory activity of curcumin: A component of tumeric (*Curcuma longa*). *J Alternat Complement Med* 2003;9:161.
17. Miquel J, Bernd A, Sempere JM, et al. The curcuma antioxidants: Pharmacological effects and prospects for future clinical use. A review. *Arch Gerontol Geriatr* 2002;34:37.
18. Chan MM, Huang HI, Fenton MR, et al. *In vivo* inhibition of nitric oxide synthase gene expression by curcumin, a cancer preventive natural product with anti-inflammatory properties. *Biochem Pharmacol* 1998;55:1955.
19. Huang TS, Lee SC, Lin JK. Suppression of c-Jun/AP-1 activation by an inhibitor of tumor promotion in mouse fibroblast cells. *Proc Natl Acad Sci USA* 1991;88:5292.
20. Bierhaus A, Zhang Y, Quehenberger P, et al. The dietary pigment curcumin reduces endothelial tissue factor gene expression by inhibiting binding of AP-1 to the DNA and activation of NF- κ B. *Thromb Haemost* 1997;77:772.
21. Abe Y, Hashimoto S, Horie T. Curcumin inhibition of inflammatory cytokine production by human peripheral blood monocytes and alveolar macrophages. *Pharmacol Res* 1999; 39:41.
22. Jang MK, Sohn DH, Ryu JH. A curcuminoid and sesquiterpenes as inhibitors of macrophage TNF- α release from *Curcuma zedoaria*. *Planta Med* 2001;67:550.
23. Egan ME, Pearson M, Weiner SA, et al. Curcumin, a major constituent of turmeric, corrects cystic fibrosis defects. *Science* 2004;304:600.
24. Siddiqui A, Cui X, Wu R, et al. The anti-inflammatory effect of curcumin in an experimental model of sepsis is mediated by up regulation of peroxisome proliferator-activated receptor- γ . *Crit Care Med* 2006;34:1874.
25. Toda S, Miyase T, Arichi H, et al. Natural antioxidants. III. Antioxidative components isolated from rhizome of *Curcuma longa* L. *Chem Pharm Bull* 1985;33:1725.
26. Sompamit K, Kukongviriyapan U, Nakmareong S, et al. Curcumin improves vascular function and alleviates oxidative stress in non-lethal lipopolysaccharide-induced endotoxemia in mice. *Eur J Pharmacol* 2009;616:192.
27. Boyne AF, Ellman GL. A methodology for analysis of tissue sulfhydryl components. *Anal Biochem* 1972;46:639.
28. Sun Y, Oberley LW, Li Y. A simple method for clinical assay of superoxide dismutase. *Clin Chem* 1988;34:497.
29. Durak I, Yurtarslani Z, Canbolat O, et al. A methodological approach to superoxide dismutase (SOD) activity assay based on inhibition of nitroblue tetrazolium (NBT) reduction. *Clin Chim Acta* 1993;214:103.
30. Aebi H. Catalase. In: Bergmeyer U, Ed. *Methods of enzymatic analysis*. New York: Academic Press, 1974:673-7.
31. Beuge JA, Aust SD. Microsomal lipid peroxidation. *Methods Enzymol* 1978;52:302.
32. Lowry OH, Rosebrough NJ, Farr AL, et al. Protein measurements with the folin phenol reagent. *J Biol Chem* 1951;193:265.
33. Levine RL, Garland D, Oliver CN, et al. Determination of carbonyl content in oxidatively modified proteins. *Meth Enzymol* 1990;186:464.
34. Dhalla NS, Elmoselhi AB, Hata T, et al. Status of myocardial antioxidants in ischemiareperfusion injury. *Cardiovasc Res* 2000; 47:446.
35. Granger DN. Role of xanthine oxidase and granulocytes in ischemia-reperfusion injury. *Am J Physiol* 1998;255:H1269.
36. Wall J. Antioxidants in prevention of reperfusion damage of vascular endothelium. *TSMJ* 2000;1:67.
37. Anaya-Prado R, Toledo-Pereyra LH, Lentsch AB, et al. Ischemia/reperfusion injury. *J Surg Res* 2002;105:248.
38. Halliwell B, Gutteridge JM, Cross CE. Free radicals, antioxidants, and human disease: Where are we now? *J Lab Clin Med* 1992;119:598.
39. Cetin C, Kose AA, Aral E, et al. Protective effect of fucoidin (a neutrophil rolling inhibitor) on ischemia-reperfusion injury: Experimental study in rat epigastric island flaps. *Ann Plast Surg* 2001;47:540.
40. Gurlek A, Celik M, Parlakpınar H, et al. The protective effect of melatonin on ischemia reperfusion injury in the groin (inferior epigastric) flap model in rats. *J Pineal Res* 2006;40:312.
41. Da Silveira M, Yoshida WB. Ischemia and reperfusion in skeletal muscle injury mechanism and treatment perspectives. *J Vasc Br* 2004;3:367.
42. Slater TF. Free-radical mechanisms in tissue injury. *Biochem J* 1984;222:1.
43. Belkin M, LaMorte WL, Wright JG, et al. The role of leukocytes in the pathophysiology of skeletal muscle ischemic injury. *J Vasc Surg* 1989;10:14.
44. Petrasko PF, Homer-Vanniasinkam S, Walker PM. Determinants of ischemic injury to skeletal muscle. *J Vasc Surg* 1994; 19:623.
45. Ikeda K, Negishi H, Yamori Y. Antioxidant nutrients and hypoxia/ ischemia brain injury in rodents. *Toxicology* 2003; 189:55.
46. Kearns SR, Daly AF, Sheehan K, et al. D. Oral vitamin C reduces the injury to skeletal muscle caused by compartment syndrome. *J Bone Joint Surg Br* 2004;86:906.
47. Erkanli K, Kayalar N, Erkanli G, et al. Melatonin protects against ischemia/reperfusion injury in skeletal muscle. *J Pineal Res* 2005;39:238.
48. Wang WZ, Fang XH, Stephenson LL, et al. Melatonin reduces ischemia/reperfusion-induced superoxide generation in arterial wall and cell death in skeletal muscle. *J Pineal Res* 2006;41:255.
49. Hosseinzadeh H, Nassiri Asl M, Parvardeh S. The effects of carbenoxolone, a semisynthetic derivative of glycyrrhizic acid, on peripheral and central ischemia-reperfusion injuries in the skeletal muscle and hippocampus of rats. *Phytomedicine* 2005;12:632.
50. Inokuchi Y, Shimazawa M, Nakajima Y, et al. Brazilian green propolis protects against retinal damage *in vitro* and *in vivo*. *Evidence Based Complement Alternat Med* 2006;3:71.
51. Shimazawa M, Chikamatsu S, Morimoto N, et al. Neuroprotection by Brazilian green propolis against *in vitro* and *in vivo*

- ischemic neuronal damage. Evidence Based Complement Alternat Med 2005;2:201.
52. Ghoneim AI, Abdel-Naim AB, Khalifa AE, et al. Protective effects of curcumin against ischemia/reperfusion insult in rat forebrain. *Pharmacol Res* 2002;6:273.
 53. Fleming SD, Starnes BW, Kiang JG, et al. Heat stress protection against mesenteric I/R-induced alterations in intestinal mucosa in rats. *J Appl Physiol* 2002;92:2600.
 54. Shen SQ, Zhang Y, Xiang JJ, et al. Protective effect of curcumin against liver warm ischemia/reperfusion injury in rat model is associated with regulation of heat shock protein and antioxidant enzymes. *World J Gastroenterol* 2007;13:1953.
 55. Sun J, Yang D, Li S, et al. Effects of curcumin or dexamethasone on lung ischaemia-reperfusion injury in rats. *Eur Respir J* 2009; 33:398.
 56. Chen KH, Chao D, Liu CF, et al. Curcumin attenuates airway hyperactivity induced by ischemia-reperfusion of the pancreas in rats. *Transplantation Proc* 2010;42:744.
 57. Appell HJ, Glöser S, Soares JMC, et al. Structural alterations of skeletal muscle induced by ischemia and reperfusion. *Basic Appl Myol* 1999;9:263.
 58. Novelli GP, Adembri C, Gandini E, et al. Vitamin E protects human skeletal muscle from damage during surgical ischemia-reperfusion. *Am J Surg* 1996;173:206.