

Original Article

The effect of different concentrations of topical ozone administration on bone formation in orthopedically expanded suture in rats

S. Kutalmış Büyük*, Sabri İlhan Ramoğlu**, and Mehmet Fatih Sonmez***

*Department of Orthodontics, Faculty of Dentistry, Ordu University, Ordu, Turkey, **Department of Orthodontics, Faculty of Dentistry, Bezmialem Vakıf University, Istanbul, Turkey and ***Department of Histology and Embryology, Faculty of Medicine, Erciyes University, Kayseri, Turkey

Correspondence to: S. Kutalmış Büyük, Department of Orthodontics, Faculty of Dentistry, Ordu University, 52100 Ordu-Turkey. E-mail: sk_buyuk@yahoo.com

Summary

Background/objective: The aim of this study was to investigate the effects of different concentrations of ozone (O₃) therapy on bone regeneration in response to an expansion of the inter-premaxillary suture in rats.

Materials and methods: Forty-eight Wistar rats were randomly divided into four groups ($n = 12$). In groups I, II, and III, 1 ml of O₃ at 10, 25, and 40 µg/ml was injected at the premaxillary suture, respectively. In group IV (control group), 1 ml of saline solution was injected at the same point during the expansion procedure for 5 days. Bone regeneration in the suture was evaluated histomorphometrically. The area of new bone and fibrotic area, the number of osteoblasts and osteoclasts, and the amount of vascularity were measured and compared. The density of the newly formed bone in the expansion area was measured by using cone beam computed tomography. Data were analyzed using the Kruskal–Wallis one-way analysis of variance and *post hoc* Student–Newman–Keuls tests.

Results: New bone area, fibrotic area, osteoblast and osteoclast numbers, and the amount of vascularity were significantly higher in experimental groups compared with the control group ($P < 0.001$). The density of newly formed bone ($P < 0.001$), new bone formation ($P = 0.009$), number of capillaries ($P < 0.001$), number of osteoclasts ($P = 0.016$), and number of osteoblasts ($P < 0.001$) in the maxillary sutures were highest in the 25 µg/ml O₃ group compared with the other experimental groups and control group.

Conclusions/implications: The application of O₃ therapy can stimulate bone regeneration in an orthopedically expanded inter-premaxillary suture during both the expansion and retention periods.

Introduction

Rapid maxillary expansion (RME) is one of the most common treatment protocols for expanding a transversally narrow maxilla (1). This treatment protocol increases the posterior dental arch

width in the transverse dimension, ensuring the ideal size of the maxilla (2).

Despite the fact that the reasons for a post-expansion relapse are not completely understood, the quality and rigidity in the sutural region during and after expansion may affect a post-treatment

relapse (3). Potentially favourable effects of accelerating bone formation in the sutural area during and after expansion have been reported to prevent a relapse of the transversal arch width and shorten the retention period (3, 4).

Ozone (O_3) is a three-atom molecule consisting of three oxygen atoms. O_3 has been used in both an aqueous and gaseous form in medicine and dentistry (5). O_3 therapy has antimicrobial, anti-inflammatory, and wound healing effects (6–8). Additionally, O_3 therapy has immunostimulating, antihypoxic, analgesic, detoxicating, bioenergetic, and biosynthetic effects on the human body (9).

O_3 provides for the synthesis of a group of cytokines, such as leukotrienes, interleukins, and prostaglandins. O_3 influences both the cellular and humoral immune system and stimulates the proliferation of immunocompetent cells and the synthesis of immunoglobulins. O_3 also activates the function of macrophages and increases the sensitivity of microorganisms to phagocytosis. There are different ways to administer O_3 gas, including minor and major autohemotherapies, insufflations, rectally, intraarticular, and local and topical applications (10).

The aim of this study was to investigate the effects of various concentrations of O_3 therapy on bone regeneration in response to the expansion of the inter-premaxillary suture in rats. These effects were evaluated with quantitative bone histomorphometric and cone beam computed tomography (CBCT) examinations. The research hypothesis of this study states that the administration of O_3 therapy has positive effects on bone formation during the inter-premaxillary suture expansion in rats.

Materials and methods

Forty-eight 11–12 week-old Wistar albino male rats with a mean weight of 203.47 ± 9.19 g were randomly divided into four groups of 12 animals each. Ethical approval for this study was obtained from the University of Erciyes Regional Animal Research Ethics Committee (13 February 2013-13/23). All rats were kept in polycarbonate cages and subjected to a 12-hour light-dark cycle at the constant temperature of 23°C and fed a normal pellet diet (Expanded pellets, Stepfield Witham, Essex, UK) with water provided *ad libitum*.

Sutural expansion was carried out in all animals using an expansion spring made of a 0.014 inch stainless steel wire inserted in holes drilled close to the gingival margins of both upper incisors. The springs were activated to deliver a force of 50 g and were not reactivated during the expansion period (11, 12) (Figure 1A).

After the expansion period, the springs were removed and a 0.016×0.016 inch stainless steel rectangular wire was inserted into the holes between two incisors during a 10-day retention period (Figure 1B). The appliances were attached to the maxillary incisors of all animals under general anesthesia with 90 mg/kg




Figure 1. (A) Expansion appliance *in situ*. (B) Retaining wire placed between rat incisors' holes.

ketamine hydrochloride (Ketazol, Wels, Austria) and 3 mg/kg xylazine (Rompun-Bayer, Leverkusen, Germany) *via* an intraperitoneal injection.

In this histomorphometrical and CBCT study, three experimental groups were treated with different O_3 gas concentrations (group I: 10 $\mu\text{g/ml}$; group II: 25 $\mu\text{g/ml}$; and group III: 40 $\mu\text{g/ml}$). O_3 therapy was performed using an O_3 generator (Ozonosan® Dr. J. Hänslér GmbH, Iffezheim, Germany) and all O_3 therapy volumes in the experimental groups were 1 ml. The O_3 therapy injections were repeated five times during the expansion period. The control group (Group IV) received a 1 ml saline solution five times during the expansion procedure (5 days). One day after the expansion appliance placement, various oxygen- O_3 concentrations, or saline solutions, were injected into the inter-premaxillary suture with a silicon-coated micro-syringe (Hamilton injection syringe, Hamilton Company, NV, USA).

The density of the newly formed bone was measured using a CBCT. All axial images were obtained in the standard position by CBCT (NewTom 5G, QR Verona, Italy). The machine scanning time was 14–18 seconds with a limited field of view ($8\text{ cm} \times 8\text{ cm}$), a voxel size of 0.3 mm^3 , and an exposure time of 3.6 seconds. The CBCT images were transformed to the Digital Imaging and Communications in Medicine (DICOM) format, and then the SimPlant Pro 13.0 software (Materialise HQ, Leuven, Belgium) was used to perform the density measurements of the newly formed bone as a Hounsfield unit (HU) (Figure 2). CBCT measurements were taken at the end of the expansion (T1) and after the retention period (T2).

At the end of the experiment, animals were sacrificed by decapitation under intraperitoneal ketamine (75 mg/kg) + xylazine (10 mg/kg) anesthesia. After decapitation, the pre-maxillae were removed instantly and fixed in 10% formalin for 10 days. Then, the pre-maxillae were decalcified with 5% ethylenediaminetetraacetic acid for 8 weeks. During decalcification, the solution was changed twice a week. The pre-maxillae were cut perpendicularly to the sagittal plane, which was determined by two points: one at the alveolar crest and the other 4 mm apically. This plane passed through the centre of the incisor crown at its gingival portion. The decalcified pre-maxillae were rinsed under running tap water for 24 hours, followed by dehydration through a graded alcohol series. Tissues were made transparent in xylol and embedded in paraffin. Five-micrometer-thick sections were stained with hematoxylin-eosin (H&E) and Masson's trichrome, and photographs were taken with an Olympus BX-51 photomicroscope.

Histomorphometric evaluations were performed in a blind analysis by two experienced histologists, and the results were an average of the counts. Because bone formation on the surface area was




Figure 2. (A) Cone beam computed tomography images after the expansion and (B) retention period.

sometimes irregular and not suitable for quantitative measurement, bone histomorphometric measurements were performed 200 μm under the surface of the osseous palate facing the oral cavity. Three non-sequential histological sections were analysed for each animal. The associated analysed parameters were the new mineralised area, the fibrosis area, the number of blood vessels, and the number of osteoblast cells within the premaxillary suture (Figures 3A–C). The differences between the groups were analysed using the Image J program (US National Institutes of Health, Bethesda, MA, USA).

Cathepsin K expression was detected immunohistochemically using a goat polyclonal antibody (sc-6507; Santa Cruz Biotechnology, CA, USA) and the streptavidin-biotin peroxidase technique. The procedure was performed under identical conditions for all sections. Paraffin sections (5 μm) were deparaffinised in xylene. The sections were rehydrated, rinsed in deionised water, and antigen retrieval was carried out by microwave treatment in a 0.01 M sodium citrate buffer (pH = 6.0) at 95°C for 5 minutes. The slides were then cooled rapidly at room temperature for 20 minutes. The sections were washed with phosphate-buffered saline (PBS), and endogenous peroxidase activity was inhibited by 3% H_2O_2 in methanol for 10 minutes. Five percent normal goat serum was used to block non-specific staining. The histological sections were then incubated with a cathepsin K-specific polyclonal antibody at a dilution of 2.5 $\mu\text{g}/\text{ml}$ in 5% serum overnight at 4°C. After washing with PBS, sections were incubated with biotinylated secondary antibodies. The immunoreaction was then amplified with the streptavidin-avidin-peroxidase complex, and the sections were visualized using 3,3'-diaminobenzidine tetrahydrochloride (DAB) and lightly counterstained with hematoxylin. When incubation with the primary antisera was omitted, negative controls were completely unlabelled.

Three slides of each specimen were stained immunohistochemically for the osteoclast (cathepsin K positive cells) count. Osteoclasts were calculated by counting in the suture area (Figure 3D).

Statistical analysis

All data was analysed with SPSS 21 (SPSS; IBM SPSS Inc., Chicago, IL, USA). When the *P*-value was less than 0.05, the statistical test was




Figure 3. (A) Histomorphometric measurements of newly formed bone area (arrow) (μm^2). (B) Histomorphometric measurements of fibrotic area (arrow) (μm^2). (C) Osteoblast cells on the newly formed bone area (arrow) and blood vessels on the fibrotic area (asterisks) (Masson-trichrome staining). (D) Osteoclast cells on the bone surface (arrow) (immunochemical staining).

determined to be significant. To evaluate and compare the density of the newly formed bone after expansion and retention (T1–T2), the Kruskal–Wallis test was used. Histologic and immunohistochemical measurements were evaluated with the Kruskal–Wallis one-way analysis of variance (ANOVA) and *post hoc* Student–Newman–Keuls tests.

Results

In the control and experimental groups, all animals survived to the end of the study. However, one animal in group I, one in group II, and one in group IV exhibited incisor tooth fractures; therefore, those animals were excluded from the study. Instead of these rats, three new rats were included in the study.

Inter-premaxillary suture separation was successfully achieved with the expansion spring. The body weight of the rats in the control and experimental groups decreased during the expansion period, but subsequently increased during the retention period.

The median densities of the newly formed bone at T1 were 810.64, 849.69, 802.84, and 691.70 HU in groups I, II, III, and IV, respectively. There were significant differences between the groups in the density of the newly formed bone at T1 ($P < 0.001$). The differences were also statistically significant between groups I and II, I and IV, and III and IV (Table 1). Moreover, the median densities of the newly formed bone at T2 were 880.28, 971.81, 885.49, and 751.23 HU in groups I, II, III, and IV, respectively. There were also significant differences between the groups in the density of the newly formed bone at T2 ($P < 0.001$). The highest bone density value was observed in group II (971.81 HU) (Table 2).

Statistical analyses showed statistically significant differences among groups for all investigated histologic and immunohistochemical parameters. New bone area, fibrotic area, blood vessels, number of osteoblasts, and osteoclasts measurements showed statistically significant differences (Tables 3 and 4; Figure 3). For all investigated histomorphometric parameters, group II showed more

Table 1. Measurement of bone density at the end of expansion period (T1) (Hounsfield Unit).

Groups	N	Median	%25p	%75p	P
Group I	12	810.64 ^b	778.49	821.33	<0.001
Group II	12	849.69 ^a	842.40	901.87	
Group III	12	802.84 ^b	765.56	822.32	
Group IV	12	691.70 ^c	669.22	753.00	

N, sample size; group I, 10 $\mu\text{g}/\text{ml}$ O_3 ; group II, 25 $\mu\text{g}/\text{ml}$ O_3 ; group III, 40 $\mu\text{g}/\text{ml}$ O_3 ; group IV, control; *p*, represents the differences among the groups; same letters represent the similarity, while different letters represent the differences among the groups.

Table 2. Measurement of bone density at the end of retention period (T2) (Hounsfield Unit).

Groups	N	Median	%25p	%75p	P
Group I	12	880.28 ^b	813.70	929.02	<0.001
Group II	12	971.81 ^a	941.68	999.99	
Group III	12	885.49 ^b	826.39	912.27	
Group IV	12	751.23 ^c	725.09	835.43	

N, sample size; group I, 10 $\mu\text{g}/\text{ml}$ O_3 ; group II, 25 $\mu\text{g}/\text{ml}$ O_3 ; group III, 40 $\mu\text{g}/\text{ml}$ O_3 ; group IV, control; *p*, represents the differences among the groups; same letters represent the similarity, while different letters represent the differences among the groups.

Table 3. Descriptive values, Kruskal–Wallis results of histomorphometric measurements

Parameters	Groups	N	Median	%25p	%75p	P
New bone area (μm^2)	Group I	12	651098.80 ^b	623055.50	662371.20	0.009
	Group II	12	764027.00 ^a	675272.00	865265.70	
	Group III	12	598194.80 ^b	514446.00	730714.30	
	Group IV	12	484390.00 ^b	397921.70	697768.00	
Fibrotic area (μm^2)	Group I	12	732997.50 ^a	719568.20	766488.50	0.007
	Group II	12	706363.20 ^b	634819.30	812325.80	
	Group III	12	590404.50 ^c	555543.00	673277.00	
	Group IV	12	520008.50 ^c	459075.00	709677.50	
Osteoclast	Group I	12	1.00 ^b	0.67	1.50	0.016
	Group II	12	2.00 ^a	1.33	2.67	
	Group III	12	1.00 ^b	0.84	1.50	
	Group IV	12	0.50 ^b	0.33	1.17	

N, sample size; group I, 10 $\mu\text{g/ml}$ O_3 ; group II, 25 $\mu\text{g/ml}$ O_3 ; group III, 40 $\mu\text{g/ml}$ O_3 ; group IV, control; *p*, represents the differences among the groups; same letters represent the similarity, while different letters represent the differences among the groups.

Table 4. Descriptive values, one-way analysis of variances and multiple comparison results of osteoblast and blood vessels number in all groups

Parameters	Groups	N	Mean	SD	Minimum	Maximum	P
Osteoblast	Group I	12	363.03 ^a	41.31	270.00	462.00	<0.001
	Group II	12	423.97 ^b	19.95	370.00	490.00	
	Group III	12	395.03 ^b	34.52	348.00	486.00	
	Group IV	12	263.33 ^c	52.35	137.00	340.00	
Blood vessel	Group I	12	23.58 ^a	5.36	16.00	41.00	<0.001
	Group II	12	28.47 ^a	7.20	13.00	57.00	
	Group III	12	24.75 ^a	3.23	17.00	36.00	
	Group IV	12	13.31 ^b	5.41	5.00	27.00	

N, sample size; group I, 10 $\mu\text{g/ml}$ O_3 ; group II, 25 $\mu\text{g/ml}$ O_3 ; group III, 40 $\mu\text{g/ml}$ O_3 ; group IV, control; *p*, represents the differences among the groups; same letters represent the similarity, while different letters represent the differences among the groups.

positive results than groups I, III, and IV with respect to new bone formation; these measurements also revealed that bone architecture in the treatment groups was improved ($P < 0.001$).

Discussion

Researchers have performed many studies involving the application of different pharmacological and biochemical agents to increase bone formation in distraction osteogenesis and sutural expansion. However, there are few studies in the orthodontic literature on stimulating regeneration in the mid-palatal suture after expansion.

In the orthodontic literature, many studies have been performed to accelerate regeneration in the inter-premaxillary suture after expansion. Sawada and Shimizu (4) applied a single dose of transforming growth factor- β 1 for stimulating the expanding rat sutures. Furthermore, Chang *et al.* (13) evaluated the effect of an angiogenic factor on cell migration patterns and osteoblast histogenesis orthopedic expansion of the anterior maxillary suture in rats. All of these studies significantly stimulated bone regeneration in the mid-palatal/inter-premaxillary suture. Recently, Ekizer *et al.* (14, 15) investigated the effects of a light-emitting diode photobiomodulation and bone marrow-derived mesenchymal stem cells on bone formation in the inter-premaxillary suture in rats. This group found that these applications stimulated bone regeneration during both the expansion and retention periods. Bone marrow mesenchymal stem cells enhance bone formation in orthodontically expanded maxillae in rats. In the current study, we investigated the effects of different concentrations of O_3 therapy on bone regeneration in response to the expansion of

the inter-premaxillary suture in rats. We found increases in the newly formed mineralised bone area in the suture area with various O_3 concentrations, especially the concentration used in group II.

Different animal models have been described for examining bone formation on expanding the inter-premaxillary suture area. Various animals, including monkeys, dogs, cats, sheep, rabbits, and rats have been used as animal models to examine the effects of biomechanical stimulation on bone (16). Monkeys and cats have similar anatomical features regarding the maxillary sutures that are, in most respects, similar to that of man; thus, these animals have been used in many maxillary expansion studies. However, the ideal animals are rabbits and rats for maxillary expansion due to the production of a clear picture of maxillary and sutural changes under stimulation and the ease of manufacturing (17). In accordance with the literature, we used rats as the experimental animal model (11, 12, 14, 15).

In this study, the positive effects of O_3 therapy on the quantity of new bone formation during maxillary expansion were investigated using a histomorphometric and CBCT evaluation. Bone histomorphometry is a reliable histology technique commonly used in quantitative evaluations of bone remodelling in experimental studies (2–4, 18). Additionally, the density of the newly formed bone in the expansion area, both after the expansion and after the retention period, was measured with the CBCT device. CBCT measurements are reliable and accurate to a subvoxel size and can potentially be used as a quantitative orthodontic diagnostic tool (19, 20).

Currently, O_3 therapy is widely used in medicine, dentistry, and maxillofacial surgery (7, 21–23); however, its application in orthodontics is limited. Ozdemir *et al.* (24) aimed to evaluate the effect

of O₃ therapy on autogenous bone graft healing in calvarial defects. They concluded that O₃ therapy enhances new bone formation using an autogenous bone graft in the rat calvarial defect model. Another study also reported that O₃ therapy has a positive effect on bone formation in rat calvarial defects compared with controls (10).

O₃ activates a series of biological mechanisms that lead to normalising the delivery of oxygen for several days with consequent therapeutic effects (5). Additionally, in the present study, the histomorphometric results showed faster bone formation in the 25 µg/ml dose of the O₃ therapy administrated group. We think that this might be due to the fact that a low concentration of O₃ therapy stimulates the immune system, while a high concentration of O₃ therapy inhibits the immune system.

Agrillo *et al.* (25) carried out 20 tooth extractions from 15 patients with avascular bisphosphonate-related jaw osteonecrosis. They reported that O₃ was effective when used 7 days before and 7 days after the tooth extraction. In our study, O₃ therapy enhanced new bone formation in the inter-premaxillary sutural area. These healing properties are due to the reaction of O₃ with erythrocytes, leukocytes, and platelets. As a result, O₃ therapy provides greater microcirculation and improves wound healing (26).

Conclusions

The application of O₃ therapy can stimulate bone regeneration in an orthopedically expanded inter-premaxillary suture during both the expansion and retention periods. Furthermore, a 25 µg/ml concentration of O₃ therapy was the most effective dose for bone regeneration in rats. Further animal and clinical studies are required to determine any pharmaceutical benefits of O₃ therapy.

Funding

The Scientific and Technological Research Council of Turkey (TUBITAK 113S118).

Acknowledgements

The authors would also like to thank Dr Mustafa Koseahmetoglu for many helpful suggestions concerning O₃ applications.

References

- Nicholson, P. and Plint, D. (1989) A long-term study of rapid maxillary expansion and bone grafting in cleft lip and palate patients. *European Journal of Orthodontics*, 11, 186–192.
- Uysal, T., Ustul, A., Sonmez, M.F. and Ozturk, F. (2009) Stimulation of bone formation by dietary boron in an orthopedically expanded suture in rabbits. *The Angle Orthodontist*, 79, 984–990.
- Saito, S. and Shimizu, N. (1997) Stimulatory effects of low-power laser irradiation on bone regeneration in midpalatal suture during expansion in the rat. *American Journal of Orthodontics and Dentofacial Orthopedics*, 111, 525–532.
- Sawada, M. and Shimizu, N. (1996) Stimulation of bone formation in the expanding mid-palatal suture by transforming growth factor-β1 in the rat. *European Journal of Orthodontics*, 18, 169–179.
- Bocci, V.A. (2006) Scientific and medical aspects of ozone therapy. State of the art. *Archives of Medical Research*, 37, 425–435.
- Ripamonti, C.I., Cislighi, E., Mariani, L. and Maniezzo, M. (2011) Efficacy and safety of medical ozone (O₃) delivered in oil suspension applications for the treatment of osteonecrosis of the jaw in patients with bone metastases treated with bisphosphonates: preliminary results of a phase I–II study. *Oral Oncology*, 47, 185–190.
- Bocci, V. (2004) Ozone as Janus: this controversial gas can be either toxic or medically useful. *Mediators of Inflammation*, 13, 3–11.

- Grigor'ian, A., Grigor'iants, L. and Guchetl, M. (2007) Experimental-morphological study of the anti-inflammatory action of ozone-perfluorane complex application [Article in Russian]. *Stomatologiya (Mosk)*, 87, 4–9.
- Seidler, V., Linetskiy, I., Hubáková, H., Stankova, H., Smucler, R. and Mazanek, J. (2008) Ozone and its usage in general medicine and dentistry. A review article. *Prague Medical Report*, 109, 5–13.
- Kazancioglu, H.O., Ezirganli, S. and Aydin, M.S. (2013) Effects of laser and ozone therapies on bone healing in the calvarial defects. *Journal of Craniofacial Surgery*, 24, 2141–2146.
- Uysal, T., Amasyali, M., Olmez, H., Enhos, S., Karslioglu, Y. and Gunhan, O. (2011) Effect of vitamin C on bone formation in the expanded inter-premaxillary suture. Early bone changes. *Journal of Orofacial Orthopedics*, 72, 290–300.
- Uysal, T., Gorgulu, S., Yagci, A., Karslioglu, Y., Gunhan, O. and Sagdic, D. (2011) Effect of resveratrol on bone formation in the expanded inter-premaxillary suture: early bone changes. *Orthodontics & Craniofacial Research*, 14, 80–87.
- Chang, H.N., Garetto, L.P., Katona, T.R., Potter, R.H. and Eugene Roberts, W. (1996) Angiogenic induction and cell migration in an orthopaedically expanded maxillary suture in the rat. *Archives of Oral Biology*, 41, 985–994.
- Ekizer, A., Uysal, T., Güray, E. and Yüksel, Y. (2013) Light-emitting diode photobiomodulation: effect on bone formation in orthopedically expanded suture in rats—early bone changes. *Lasers in Medical Science*, 28, 1263–1270.
- Ekizer, A., Yalvac, M.E., Uysal, T., Sonmez, M.F. and Sahin, F. (2015) Bone marrow mesenchymal stem cells enhance bone formation in orthodontically expanded maxillae in rats. *The Angle Orthodontist*, 85, 394–399.
- Swennen, G., Dempf, R. and Schliephake, H. (2002) Cranio-facial distraction osteogenesis: a review of the literature. Part II: experimental studies. *International Journal of Oral and Maxillofacial Surgery*, 31, 123–135.
- Storey, E. (1973) Tissue response to the movement of bones. *American Journal of Orthodontics*, 64, 229–247.
- Uysal, T., Amasyali, M., Enhos, S., Sonmez, M.F. and Sagdic, D. (2009) Effect of ED-71, a new active vitamin D analog, on bone formation in an orthopedically expanded suture in rats. A histomorphometric study. *European Journal of Dentistry*, 3, 165–172.
- Gribel, B.F., Gribel, M.N., Frazão, D.C., McNamara, J.A., Jr and Manzi, F.R. (2011) Accuracy and reliability of craniometric measurements on lateral cephalometry and 3D measurements on CBCT scans. *The Angle Orthodontist*, 81, 26–35.
- Celikoglu, M., Halicioglu, K., Buyuk, S.K., Sekerci, A.E. and Ucar, F.I. (2013) Condylar and ramal vertical asymmetry in adolescent patients with cleft lip and palate evaluated with cone-beam computed tomography. *American Journal of Orthodontics and Dentofacial Orthopedics*, 144, 691–697.
- Martínez-Sánchez, G., Al-Dalain, S.M., Menéndez, S., Re, L., Giuliani, A., Candelario-Jalil, E., Alvarez, H., Fernández-Montequín, J.I. and León, O.S. (2005) Therapeutic efficacy of ozone in patients with diabetic foot. *European Journal of Pharmacology*, 523, 151–161.
- Magni, E., Ferrari, M., Pacacchini, F., Hickel, R. and Ilie, N. (2010) Influence of ozone application on the repair strength of silorane-based and ormocer-based composites. *American Journal of Dentistry*, 23, 260–264.
- Stübinger, S., Sader, R. and Filippi, A. (2006) The use of ozone in dentistry and maxillofacial surgery: a review. *Quintessence International*, 37, 353–359.
- Ozdemir, H., Tokar, H., Balci, H. and Ozer, H. (2013) Effect of ozone therapy on autogenous bone graft healing in calvarial defects: a histologic and histometric study in rats. *Journal of Periodontal Research*, 48, 722–726.
- Agrillo, A., Ungari, C., Filiaci, F., Priore, P. and Iannetti, G. (2007) Ozone therapy in the treatment of avascular bisphosphonate-related jaw osteonecrosis. *Journal of Craniofacial Surgery*, 18, 1071–1075.
- Azarpazhooh, A. and Limeback, H. (2008) The application of ozone in dentistry: a systematic review of literature. *Journal of Dentistry*, 36, 104–116.