



# Association between food insecurity and depression among older adults from low- and middle-income countries

Lee Smith<sup>1</sup> | Jae Il Shin<sup>2</sup> | Daragh McDermott<sup>3</sup> | Louis Jacob<sup>4,5</sup> |  
 Yvonne Barnett<sup>6</sup> | Guillermo F. López-Sánchez<sup>7</sup> | Nicola Veronese<sup>8</sup> |  
 Lin Yang<sup>9</sup> | Pinar Soysal<sup>10</sup> | Hans Oh<sup>11</sup> | Igor Grabovac<sup>12</sup> | Ai Koyanagi<sup>4,13</sup>

<sup>1</sup>The Cambridge Centre for Sport and Exercise Sciences, Anglia Ruskin University, Cambridge, UK

<sup>2</sup>Department of Pediatrics, Yonsei University College of Medicine, Seoul, South Korea

<sup>3</sup>School of Psychology & Sport Science, Anglia Ruskin University, Cambridge, UK

<sup>4</sup>Research and Development Unit, Parc Sanitari Sant Joan de Déu, CIBERSAM, Barcelona, Spain

<sup>5</sup>Faculty of Medicine, University of Versailles Saint-Quentin-en-Yvelines, Montigny-le-Bretonneux, France

<sup>6</sup>Anglia Ruskin University, Cambridge, UK

<sup>7</sup>School of Medicine, Vision and Eye Research Institute, Faculty of Health, Education, Medicine and Social Care, Anglia Ruskin University-Cambridge Campus, Cambridge, UK

<sup>8</sup>Geriatric Unit, Department of Internal Medicine and Geriatrics, University of Palermo, Palermo, Italy

<sup>9</sup>Department of Cancer Epidemiology and Prevention Research and Departments of Oncology and Prevention Research, Alberta Health Services, University of Calgary, Calgary, Canada

<sup>10</sup>Faculty of Medicine, Bezmiâlem Vakıf University, Istanbul, Turkey

<sup>11</sup>Suzanne Dworak Peck School of Social Work, University of Southern California, Los Angeles, California, USA

<sup>12</sup>Department of Social and Preventive Medicine, Centre for Public Health, Medical University of Vienna, Vienna, Austria

<sup>13</sup>ICREA, Barcelona, Spain

## Correspondence

Lee Smith, The Cambridge Centre for Sport and Exercise Sciences, Anglia Ruskin University, Cambridge, UK, CB1 1PT.

Email: [Lee.Smith@aru.ac.uk](mailto:Lee.Smith@aru.ac.uk)

## Abstract

**Background:** To examine the association between self-reported food insecurity and depression in 34,129 individuals aged  $\geq 50$  years from six low- and middle-income countries (LMICs) (China, Ghana, India, Mexico, Russia, and South Africa).

**Methods:** Cross-sectional, community-based, nationally representative data from the WHO Study on global AGEing and adult health (SAGE) were analyzed. Self-reported past 12-month food insecurity was assessed with two questions on frequency of eating less and hunger due to lack of food. Questions based on the World Mental Health Survey version of the Composite International Diagnostic Interview were used for the endorsement of past 12-month DSM-IV depression. Multivariable logistic regression analysis and meta-analysis were conducted to assess associations.

**Results:** In total, 34,129 individuals aged  $\geq 50$  years [mean (*SD*) age, 62.4 (16.0) years; 52.1% females] were included in the analysis. Overall, the prevalence of moderate and severe food insecurity was 6.7% and 5.1%, respectively, while the prevalence of depression was 6.0%. Meta-analyses based on countrywise estimates showed that overall, moderate food insecurity (vs. no food insecurity) is associated with a nonsignificant 1.69 (95% confidence interval [CI] = 0.82–3.48) times higher odds for depression, while severe food insecurity is significantly associated with 2.43 (95% CI = 1.65–3.57) times higher odds for depression.

**Conclusions:** In this large representative sample of older adults from six LMICs, those with severe food insecurity were over two times more likely to suffer from depression (compared with no food insecurity). Utilizing lay health counselors and psychological interventions may be effective mechanisms to reduce depression among food-insecure populations. Interventions to address food insecurity (e.g., supplemental nutrition programs) may reduce depression at the population level but future longitudinal studies are warranted.

## KEY WORDS

depression, food insecurity, low- and middle-income countries, older adults

## 1 | INTRODUCTION

Depression is characterized by persistent sadness and a lack of interest or pleasure in previously rewarding or enjoyable activities (World Health Organization, 2020a). Globally, depression is a common mental disorder affecting more than 264 million people (World Health Organization, 2020b). Importantly, the Global Burden of Disease Study 2010 demonstrated that depression is responsible for 40.5% of total disability-adjusted life years caused by mental and substance use disorders (Whiteford et al., 2015). Depression may be disproportionately affecting low- and middle-income countries (LMICs), as these countries bear more than 80% of the global years lost to disability due to depression, while the majority of people with depression in LMICs do not receive appropriate treatment (Jacob et al., 2019). Furthermore, depression in late life is of particular importance from a public health and clinical point of view as its prevalence is high (approximately 7%) (World Health Organization, 2017), while it is associated with greater risk of morbidity and suicide, poorer cognitive, physical, and social functioning, and increased levels of self-neglect, all of which may result in premature mortality (Fiske et al., 2009; Yohannes & Baldwin, 2008). It is, therefore, important to identify correlates of depression among the older population in LMICs to inform targeted prevention efforts.

To date, several correlates of depression have been identified in such settings, including, for example, female gender (Tomlinson et al., 2009), lack of social support (Cheng et al., 2014), low self-esteem (Azizi et al., 2013), stressful life events (Tao et al., 2011), family history of psychiatric disorders (Chin et al., 2016), chronic conditions (Das et al., 2013), and low socioeconomic status (SES) (Fernández-Niño et al., 2014). With respect to low SES, several studies have found that indicators of SES are associated with depression, and that income alone is not necessarily an adequate measure of SES, given the multidimensional nature of poverty (Alkire & Foster, 2011; Alkire et al., 2015). In this regard, one example was the study of Zimmerman and Katon (2005) which suggests that employment status and the ratio of debt-to-assets (but not income) were causally related to depression. As such, food insecurity (i.e., state of being without reliable access to a sufficient quantity of affordable, nutritious food) can be considered to be another dimension of SES linked to depression. Importantly, food insecurity may be related to depression via poor nutrition or increased feelings of shame, which are factors that can increase the risk for depression (Gilbert, 2000; Li et al., 2017).

One study carried out in Mexico on 33,011 participants with a mean age of 44 years found that participants with moderate food insecurity (odds ratio [OR] = 1.45; 95% confidence interval [CI] = 1.28–1.64) and severe food insecurity (OR = 2.04; 95% CI = 1.76–2.37) were more likely to suffer from depression as compared with participants with mild food insecurity (Kolovos et al., 2020). Moreover, another study carried out in South Africa on 8801 participants, found that those who were depression-free at baseline, and resided in a food-insecure hotspot community had a significantly higher incidence of depression (adjusted relative

risk = 1.11; 95% CI = 1.01–1.22) compared with those who resided in a food-secure hotspot (Tomita et al., 2020). However, apart from these studies, there are no other studies from LMICs that used nationally representative data sets of the general population. Other studies from LMICs that have focused on the relationship between food insecurity and depression have utilized specific populations such as mothers who recently gave birth and people living with HIV, and thus, generalizability is severely limited (Ayano et al., 2020; Dewing et al., 2013; Khoshgoo et al., 2020; Natamba et al., 2017; Tuthill et al., 2020).

Moreover, there is very limited information on this topic among older adults from this setting (Isaura et al., 2019; Mesbah et al., 2020). Thus, it is clear that more research is needed in large representative samples of older adults from diverse LMICs to further elucidate on the association between food insecurity and depression, especially given the high prevalence of both conditions in these countries (United Nations, 2019). Furthermore, multicountry studies are needed as such studies allow exploration of associations between depression and food insecurity irrespective of national policies and available facilities, and at the same time allow comparison between countries to understand whether associations are context-specific.

Therefore, the aim of the present study was to examine the association between food insecurity and depression in 34,129 individuals aged ≥50 years from six LMICs (China, Ghana, India, Mexico, Russia, and South Africa). It was hypothesized that those who experience food insecurity will have higher odds for depression compared with those who do not suffer from food insecurity.

## 2 | METHODS

### 2.1 | The survey

Data from the Study on global AGEing and adult health (SAGE) were analyzed. These data are publicly available through <http://www.who.int/healthinfo/sage/en/>. This survey was undertaken in China, Ghana, India, Mexico, Russia, and South Africa between 2007 and 2010. Based on the World Bank classification at the time of the survey, Ghana was the only low-income country, and China and India were lower middle-income countries although China became an upper middle-income country in 2010. The remaining countries were upper middle-income countries.

Details of the survey methodology have been published previously (Kowal et al., 2012). Briefly, to obtain nationally representative samples, a multistage clustered sampling design method was used. The sample consisted of adults aged ≥18 years with oversampling of those aged ≥50 years. Trained interviewers conducted face-to-face interviews using a standard questionnaire. Standard translation procedures were undertaken to ensure comparability between countries. The survey response rates were China, 93%; Ghana, 81%; India, 68%; Mexico, 53%; Russia, 83%; and South Africa, 75%. Sampling weights were constructed to adjust for the population structure as reported by the United Nations Statistical

**TABLE 1** Questions and answer options used for endorsement of DSM-IV depression

1. At least one of the two following symptoms in the last 12 months:
(a) A period, lasting several days, of feeling sad, empty, or depressed
(b) A period lasting several days with a loss of interest in most things the participant usually enjoys such as personal relationships, work, or hobbies/recreation
AND
2. The period of sadness/loss of interest/low energy lasted for more than two weeks and was most of the day and nearly every day
AND
3. Five or more of the following symptoms:
(a) Loss of appetite
(b) Insomnia (problems falling asleep or waking up too early)
(c) Decreased energy or tiredness all the time
(d) Slowing down in moving around or restless/jittery
(e) Negative feelings/loss of confidence or frequent feelings of hopelessness
(f) Slowed thinking or difficulties concentrating (e.g., listening to others, working, watching TV, listening to the radio)
(g) Thoughts of death, wishes of own death or suicide attempt
(h) Feelings of sadness, emptiness, or depression lasting several days
(i) Anhedonia: loss of interest in things the participant usually enjoys

Division. Ethical approval was obtained from the WHO Ethical Review Committee and local ethics research review boards. Written informed consent was obtained from all participants.

## 2.2 | Depression

Questions based on the World Mental Health Survey version of the Composite International Diagnostic Interview (Kessler & Üstün, 2004) were used for the endorsement of past 12-month DSM-IV depression using the same algorithm in previous SAGE studies (Garin et al., 2016; Koyanagi et al., 2014) (details provided in Table 1).

## 2.3 | Food insecurity

Food insecurity was defined with the use of the two following questions: "In the last 12 months, how often did you ever eat less than you felt you should because there wasn't enough food?" and "In the last 12 months, were you ever hungry, but didn't eat because you couldn't afford enough food?" Both of these questions had as answer options: every month (coded = 1); almost every month (coded = 2); some months, but not every month (coded = 3); only in 1 or 2 months (coded = 4); never (coded = 5). These items were adapted from similar items in food security questionnaires such as the US Household Food Security Survey Module and National Health and Nutrition Examination Survey (NHANES) Food Security module. As in previous SAGE studies, those who answered 1 through 3 to both

questions or answered 1 to either item were categorized as severely food-insecure. Those who did not meet the criteria for severe food insecurity but answered 2 through 4 for either question were coded as moderately food-insecure. Those who answered 5 to both items were categorized as food-secure (Koyanagi et al., 2019; Schrock et al., 2017).

## 2.4 | Control variables

The control variables were selected based on past literature (Montgomery et al., 2017), and included age, sex, wealth quintiles based on income, highest level of education achieved (primary, secondary, tertiary), physical activity, and smoking (never, current, past). Levels of physical activity were assessed with the Global Physical Activity Questionnaire and were classified as low, moderate, and high based on conventional cut-offs (Bull et al., 2009).

## 2.5 | Statistical analysis

The statistical analysis was performed with Stata 14.1 (Stata Corp LP). The analysis was restricted to those aged  $\geq 50$  years. We conducted countrywise multivariable logistic regression analysis to assess the association between food insecurity (exposure) and depression (outcome). The regression analysis was adjusted for age, sex, wealth, education, physical activity, and smoking. Furthermore, to assess the between-country heterogeneity that may exist in the association between food insecurity and depression, we calculated Higgins's  $I^2$  based on estimates from each country. Higgins's  $I^2$  represents the degree of heterogeneity that is not explained by sampling error with a value of  $<40\%$  often considered as negligible and  $40\%-60\%$  as moderate heterogeneity (Higgins & Thompson, 2002). A pooled estimate was obtained by random-effect meta-analysis. All variables were included in the models as categorical variables with the exception of age (continuous variable). The sample weighting and the complex study design were taken into account in the analyses. Results from the regression analyses are presented as ORs with 95% CIs. The level of statistical significance was set at  $p < .05$ .

## 3 | RESULTS

A total of 34,129 individuals aged  $\geq 50$  years [mean ( $SD$ ) age, 62.4 (16.0) years; 52.1% females] were included in the current analysis. The sample size by country was as follows: 13,175 (China); 4305 (Ghana); 6560 (India); 2313 (Mexico); 3938 (Russia); 3838 (South Africa). Overall, the prevalence of moderate and severe food insecurity was 6.7% and 5.1%, respectively, while the prevalence of depression was 6.0%. The sample characteristics by country are shown in Table 2. The prevalence of depression ranged from 1.1% (China) to 12.9% (India), while the prevalence of severe food insecurity ranged from 0.3% (China) to 21.5% (South Africa). In all

Characteristic	China	Ghana	India	Mexico	Russia	South Africa
Depression						
No	98.9	92.8	87.1	89.2	96.5	97.0
Yes	1.1	7.2	12.9	10.8	3.5	3.0
Food insecurity						
None	98.8	55.4	81.5	64.9	86.1	67.6
Moderate	0.9	23.5	11.1	15.3	7.7	10.9
Severity	0.3	21.0	7.5	19.7	6.2	21.5
Age (years), mean (SD)	62.6 (16.7)	64.4 (19.9)	61.5 (13.7)	63.0 (18.9)	63.9 (15.4)	61.6 (18.4)
Sex						
Male	49.8	52.4	51.0	46.8	38.9	44.1
Female	50.2	47.6	49.0	53.2	61.1	55.9
Education						
Primary	63.0	75.3	76.1	79.6	7.5	71.4
Secondary	32.5	21.1	18.8	12.3	74.2	22.8
Tertiary	4.5	3.6	5.1	8.1	18.2	5.7
Physical activity						
High	43.6	61.5	52.2	40.0	57.4	27.8
Moderate	27.5	12.5	22.9	22.4	15.8	12.4
Low	28.8	26.0	25.0	37.6	26.8	59.8
Smoking						
Never	64.1	75.1	45.3	60.7	69.6	66.8
Current	29.3	10.7	50.0	20.3	21.3	23.8
Former	6.6	14.2	4.7	19.1	9.0	9.4

Note: Data are presented as % unless otherwise stated.

countries, the prevalence of depression was highest among those with severe food insecurity with the exception of Ghana where the prevalence was highest among those with moderate food insecurity (Figure 1). The countrywise association between moderate food insecurity (vs. no food insecurity) is shown in Figure 2. Overall, moderate food insecurity was not significantly associated with depression ( $OR = 1.69$ ;  $95\% CI = 0.82-3.48$ ) with a high level of between-country heterogeneity ( $I^2 = 85.2\%$ ). In terms of severe food insecurity (vs. no food insecurity), overall, the OR (95% CI) was 2.43 (1.65-3.57) with a low level of between-country heterogeneity ( $I^2 = 31.0\%$ ) (Figure 3).


## 4 | DISCUSSION

In this large representative sample of older adults from multiple LMICs, it was found that those who suffer from severe food insecurity compared with no food insecurity were over two times more likely to have depression. Furthermore, overall, moderate food insecurity was not associated with depression. To the best of our knowledge, this is the first study on food insecurity and depression among older adults using nationally representative data from LMICs, while it is also the first multicountry study on this topic.

**TABLE 2** Sample characteristics by country

The present findings both support and add to previous literature. They support previous literature through providing further evidence for an association between food insecurity and a higher odds of depression in LMICs (Dewing et al., 2013; Kolovos et al., 2020), and add to previous literature through showing that an association exists in a large sample of older adults from multiple LMICs, which broadly represent different geographical locations and levels of socio-economic and demographic transition.

Food insecurity may be associated with higher odds of depression through several mechanisms. First, food insecurity results in poor nutrition as when food becomes scarce, people tend to change their dietary habits to those which are cheaper but less nutritious (e.g., high fat and carbohydrates, low vitamins and micronutrients). In turn, poor nutrition has been found to be associated with a higher risk of depression (Li et al., 2017; Rao et al., 2008; Vafaei et al., 2013). Malnutrition over time can result in poor mental health; notably, antioxidants (e.g., vitamin C, vitamin E, and other carotenoids compounds) have beneficial protective effects against depression, and their deficiencies may contribute to depression (Li et al., 2017). Second, food insecurity may result in higher levels of stress (e.g., stress in relation to feeding oneself or family) and stress is associated with higher levels of depression, and this can be due to impaired hypothalamic-pituitary-adrenal axis function




**FIGURE 1** Prevalence of depression by food insecurity (FI) status


(Mayo Clinic, 2020). Indeed, low SES/poverty has been associated with higher levels of stress hormones (e.g., cortisol, epinephrine) (Cohen et al., 2006). Third, food insecurity likely increases feelings of shame, and shame may also be associated with a higher risk of depression (Gilbert, 2000). Finally, depression has been found to influence the individual's ability to obtain and retain employment, particularly for individuals with lower education and thus depression per se may increase one's odds of suffering from food insecurity (Kolovos et al., 2020). Indeed, the relationship is likely bidirectional, which has been debated in terms of social causation and social

selection/drift (Dohrenwend et al., 1992). In other words, social and economic conditions of poverty (such as food insecurity) increase the risk for mental illness (social causation), while people with mental illnesses can "drift" into poverty (due to disability, stigma, healthcare costs) (Lund & Cois, 2018). Our findings are likely to include all the aforementioned pathways.

Findings from the present study and others suggest that food insecurity may be a risk factor for depression, and that intervention and policy are needed in LMICs to reduce levels of depression among those who suffer from food insecurity. First, in high-income



**FIGURE 2** Countrywise association between moderate food insecurity (vs. no food insecurity) estimated by multivariable logistic regression. Models are adjusted for age, sex, wealth, education, physical activity, and smoking. The overall estimate was obtained by meta-analysis with random effects. CI, confidence interval; OR, odds ratio



**FIGURE 3** Countrywise association between severe food insecurity (vs. no food insecurity) estimated by multivariable logistic regression. Models are adjusted for age, sex, wealth, education, physical activity, and smoking. The overall estimate was obtained by meta-analysis with random effects. CI, confidence interval; OR, odds ratio

countries, supplemental nutrition programs have been shown to decrease food insecurity (Gundersen et al., 2018) and improve health. Such programs have the potential to even address severe food insecurity (Keith-Jennings et al., 2019). Importantly, by addressing food insecurity, there will likely be a reduction in the associated negative outcomes (e.g., poor nutrition, shame, stress) that can increase the risk for depression. However, whether or not supplemental nutrition programs work in LMICs remains to be seen. Second, countries can increase access to affordable mental health care through the use of lay health counselors, a form of task shifting already shown to be effective in the treatment of common mental disorders in LMICs (Reynolds et al., 2012). Moreover, literature suggests that psychological interventions, such as cognitive behavior therapy, can effectively reduce symptoms of depression in LMICs (Morina et al., 2017). Furthermore, if food insecurity can be proven to be a risk factor for depression in LMICs in future longitudinal and intervention studies, alleviating poverty and hunger or poor nutrition through, for example, acquisition of agricultural input, may have a positive impact on individual and population mental well-being in this setting.

It is important to consider the present findings in light of the COVID-19 pandemic. Social distancing measures implemented to reduce and slow the transmission of SARS-CoV-2 have resulted in the closure of workplaces, an increase in unemployment, and reduced earnings (International Labour Organization, 2020). This has likely increased the proportion of the global population experiencing food insecurity and subsequent depression. Implementing measures to address food insecurity and depression, such as those discussed above, may now be of high public health importance.

The large representative sample of older adults from multiple LMICs, which collectively comprise nearly half of the worldwide population (Kowal et al., 2012), is a clear strength of the present study. However, it is important to interpret findings in light of the study limitations. First, the present study was cross-sectional in nature and thus it is not possible to determine whether food insecurity leads to depression or whether depression leads to food

insecurity. As previously discussed, the association is likely to be bidirectional. Second, our measure of food insecurity was based on two questions and did not constitute a comprehensive food insecurity measure. Third, both the exposure variable (food insecurity) and the outcome variable (depression) were self-reported, potentially introducing self-report and recall bias into the findings. Finally, although we did adjust for levels of wealth in our study, we cannot preclude the possibility that food insecurity may be a marker of other types of material deprivation (e.g., clothing, treatment for illness), which may not be reflected in our measure of wealth.

## 5 | CONCLUSIONS

In this large representative sample of older adults from six LMICs, it was found that those with severe food insecurity compared with no food insecurity were over two times more likely to suffer from depression. Utilizing lay health counselors and psychological interventions may be effective mechanisms to reduce depression among food insecure populations. Furthermore, future studies should assess whether alleviating food insecurity can lead to a lower risk of depression in LMICs.

## ACKNOWLEDGMENTS

This paper uses data from WHO's Study on global AGEing and adult health (SAGE). SAGE is supported by the US National Institute on Aging through Interagency Agreements OGHA 04034785, YA1323-08-CN-0020, Y1-AG-1005-01 and through research grants R01-AG034479 and R21-AG034263.

## CONFLICT OF INTERESTS

The authors declare that there are no conflict of interests.

## AUTHOR CONTRIBUTORS

All authors have made a substantial, direct, and intellectual contribution to the work. All authors have read and approved the final

version of the manuscript, and agree with the order of presentation of the authors.

## DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

## ORCID

Louis Jacob  <https://orcid.org/0000-0003-1071-1239>  
Guillermo F. López-Sánchez  <https://orcid.org/0000-0002-9897-5273>  
Lin Yang  <https://orcid.org/0000-0002-1698-6666>

## REFERENCES

- Alkire, S., & Foster, J. (2011). Counting and multidimensional poverty measurement. *Journal of Public Economics*, 95(7–8), 476–487.
- Alkire, S., Roche, J. M., Ballon, P., Foster, J., Santos, M. E., & Seth, S. (2015). *Multidimensional poverty measurement and analysis*. Oxford University Press.
- Ayano, G., Tsegay, L., & Solomon, M. (2020). Food insecurity and the risk of depression in people living with HIV/AIDS: A systematic review and meta-analysis. *AIDS Research and Therapy*, 17(1), 1–11.
- Azizi, M., Khamseh, F., Rahimi, A., & Barati, M. (2013). The relationship between self-esteem and depression in nursing students of a selected medical university in Tehran. *Iranian Journal of Psychiatric Nursing*, 1(1), 28–34.
- Bull, F. C., Maslin, T. S., & Armstrong, T. (2009). Global physical activity questionnaire (GPAQ): Nine country reliability and validity study. *Journal of Physical Activity and Health*, 6(6), 790–804.
- Cheng, Y., Li, X., Lou, C., Sonenstein, F. L., Kalamar, A., Jejeebhoy, S., Delany-Moretlwe, S., Brahmbhatt, H., Olumide, A. O., & Ojengbede, O. (2014). The association between social support and mental health among vulnerable adolescents in five cities: Findings from the study of the well-being of adolescents in vulnerable environments. *Journal of Adolescent Health*, 55(6), S31–S38.
- Chin, W., Wan, E. Y. F., Choi, E. P. H., Chan, K. T. Y., & Lam, C. L. K. (2016). The 12-month incidence and predictors of PHQ-9-screened depressive symptoms in Chinese primary care patients. *The Annals of Family Medicine*, 14(1), 47–53.
- Cohen, S., Doyle, W. J., & Baum, A. (2006). Socioeconomic status is associated with stress hormones. *Psychosomatic Medicine*, 68(3), 414–420.
- Das, R., Singh, O., Thakurta, R. G., Khandakar, M. R., Ali, S. N., Mallick, A. K., Roy, P., & Bhattacharrya, A. K. (2013). Prevalence of depression in patients with type II diabetes mellitus and its impact on quality of life. *Indian Journal of Psychological Medicine*, 35(3), 284–289.
- Dewing, S., Tomlinson, M., le Roux, I. M., Chopra, M., & Tsai, A. C. (2013). Food insecurity and its association with co-occurring postnatal depression, hazardous drinking, and suicidality among women in peri-urban South Africa. *Journal of Affective Disorders*, 150(2), 460–465.
- Dohrenwend, B., Levav, I., Shrout, P., Schwartz, S., Naveh, G., Link, B., Skodol, A., & Stueve, A. (1992). Socioeconomic status and psychiatric disorders: The causation-selection issue. *Science*, 255(5047), 946–952.
- Fernández-Niño, J. A., Manrique-Espinoza, B. S., Bojorquez-Chapela, I., & Salinas-Rodríguez, A. (2014). Income inequality, socioeconomic deprivation and depressive symptoms among older adults in Mexico. *PLOS One*, 9(9), e108127.
- Fiske, A., Wetherell, J. L., & Gatz, M. (2009). Depression in older adults. *Annual Review of Clinical Psychology*, 5, 363–389.
- Garin, N., Koyanagi, A., Chatterji, S., Tyrovolas, S., Olaya, B., Leonardi, M., Lara, E., Koskinen, S., Tobiasz-Adamczyk, B., Ayuso-Mateos, J. L., & Haro, J. M. (2016). Global multimorbidity patterns: A cross-sectional, population-based, multi-country study. *Journals of Gerontology Series A: Biomedical Sciences and Medical Sciences*, 71(2), 205–214.
- Gilbert, P. (2000). The relationship of shame, social anxiety and depression: The role of the evaluation of social rank. *Clinical Psychology & Psychotherapy*, 7(3), 174–189.
- Gundersen, C., Kreider, B., & Pepper, J. V. (2018). Reconstructing the supplemental nutrition assistance program to more effectively alleviate food insecurity in the United States. *RSF: The Russell Sage Foundation Journal of the Social Sciences*, 4(2), 113–130.
- Higgins, J. P., & Thompson, S. G. (2002). Quantifying heterogeneity in a meta-analysis. *Statistics in Medicine*, 21(11), 1539–1558.
- International Labour Organization. (2020). COVID-19 and the world of work. Updated Estimates and Analysis. [https://www.ilo.org/wcmsp5/groups/public/@dgreports/@dcomm/documents/briefingnote/wcms\\_767028.pdf](https://www.ilo.org/wcmsp5/groups/public/@dgreports/@dcomm/documents/briefingnote/wcms_767028.pdf)
- Isaura, E. R., Chen, Y., Adi, A. C., Fan, H., Li, C., & Yang, S. (2019). Association between depressive symptoms and food insecurity among Indonesian adults: Results from the 2007–2014 Indonesia Family Life Survey. *Nutrients*, 11(12), 3026.
- Jacob, L., Pizzol, D., Veronese, N., Stubbs, B., & Koyanagi, A. (2019). Physical injury and depression in six low-and middle-income countries: A nationally representative study. *Journal of Affective Disorders*, 248, 99–107.
- Keith-Jennings, B., Llobrener, J., & Dean, S. (2019). Links of the supplemental nutrition assistance program with food insecurity, poverty, and health: Evidence and potential. *American Journal of Public Health*, 109(12), 1636–1640.
- Kessler, R. C., & Üstün, T. B. (2004). The World Mental Health (WMH) survey initiative version of the World Health Organization (WHO) Composite International Diagnostic Interview (CIDI). *International Journal of Methods in Psychiatric Research*, 13(2), 93–121.
- Khoshgoo, M., Eslami, O., Al-Hosseini, M. K., & Shidfar, F. (2020). The relationship between household food insecurity and depressive symptoms among pregnant women: A cross sectional study. *Iranian Journal of Psychiatry*, 15(2), 126.
- Kolovos, S., Zavala, G. A., Leijen, A. S., Melgar-Quiñonez, H., & van Tulder, M. (2020). Household food insecurity is associated with depressive symptoms: Results from a Mexican population-based survey. *Food Security*, 12, 407–416.
- Kowal, P., Chatterji, S., Naidoo, N., Biritwum, R., Fan, W., Lopez Ridaura, R., & Williams, S. (2012). Data resource profile: The World Health Organization Study on global AGEing and adult health (SAGE). *International Journal of Epidemiology*, 41(6), pp. 1639–1649.
- Koyanagi, A., Garin, N., Olaya, B., Ayuso-Mateos, J. L., Chatterji, S., Leonardi, M., Koskinen, S., Tobiasz-Adamczyk, B., & Haro, J. M. (2014). Chronic conditions and sleep problems among adults aged 50 years or over in nine countries: A multi-country study. *PLOS One*, 9(12), e114742.
- Koyanagi, A., Veronese, N., Stubbs, B., Vancampfort, D., Stickley, A., Oh, H., Shin, J., Jackson, S., Smith, L., & Lara, E. (2019). Food insecurity is associated with mild cognitive impairment among middle-aged and older adults in South Africa: Findings from a nationally representative survey. *Nutrients*, 11(4), 749.
- Li, Y., Lv, M., Wei, Y., Sun, L., Zhang, J., Zhang, H., & Li, B. (2017). Dietary patterns and depression risk: A meta-analysis. *Psychiatry Research*, 253, 373–382.
- Lund, C., & Cois, A. (2018). Simultaneous social causation and social drift: Longitudinal analysis of depression and poverty in South Africa. *Journal of Affective Disorders*, 229, 396–402.

- Mayo Clinic. (2020). Stress management. <https://www.mayoclinic.org/healthy-lifestyle/stress-management/expert-answers/stress/faq-20058233>
- Mesbah, S. F., Sulaiman, N., Mohd Shariff, Z., & Ibrahim, Z. (2020). Does food insecurity contribute towards depression? A cross-sectional study among the urban elderly in Malaysia. *International Journal of Environmental Research and Public Health*, 17(9), 3118.
- Montgomery, J., Lu, J., Ratliff, S., & Mezuk, B. (2017). Food insecurity and depression among adults with diabetes: Results from the national health and nutrition examination survey (NHANES). *The Diabetes Educator*, 43(3), 260–271.
- Morina, N., Malek, M., Nickerson, A., & Bryant, R. A. (2017). Meta-analysis of interventions for posttraumatic stress disorder and depression in adult survivors of mass violence in low-and middle-income countries. *Depression and Anxiety*, 34(8), 679–691.
- Natamba, B. K., Mehta, S., Achan, J., Stoltzfus, R. J., Griffiths, J. K., & Young, S. L. (2017). The association between food insecurity and depressive symptoms severity among pregnant women differs by social support category: A cross-sectional study. *Maternal & Child Nutrition*, 13(3), e12351.
- Rao, T. S., Asha, M. R., Ramesh, B. N., & Rao, K. J. (2008). Understanding nutrition, depression and mental illnesses. *Indian Journal of Psychiatry*, 50(2), 77.
- Reynolds III, C. F., Cuijpers, P., Patel, V., Cohen, A., Dias, A., Chowdhary, N., Okereke, O. I., Amanda Dew, M., Anderson, S. J., Mazumdar, S., Lotrich, F., & Albert, S. M. (2012). Early intervention to reduce the global health and economic burden of major depression in older adults. *Annual Review of Public Health*, 33, 123–135.
- Schrock, J. M., McClure, H. H., Snodgrass, J. J., Liebert, M. A., Charlton, K. E., Arokiasamy, P., Naidoo, N., & Kowal, P. (2017). Food insecurity partially mediates associations between social disadvantage and body composition among older adults in India: Results from the Study on global AGEing and adult health (SAGE). *American Journal of Human Biology*, 29(6), e23033.
- Tao, M., Li, Y., Xie, D., Wang, Z., Qiu, J., Wu, W., Sun, J., Wang, Z., Tao, D., Zhao, H., Tian, T., Zhang, J., Gao, C., Niu, Q., Li, Q., Liu, S., Liu, J., Zhang, Y., He, Q., ... Gao, J. (2011). Examining the relationship between lifetime stressful life events and the onset of major depression in Chinese women. *Journal of Affective Disorders*, 135(1–3), 95–99.
- Tomita, A., Cuadros, D. F., Mabhaudhi, T., Sartorius, B., Ncama, B. P., Dangour, A. D., Tanser, F., Modi, A. T., Slotow, R., & Burns, J. K. (2020). Spatial clustering of food insecurity and its association with depression: A geospatial analysis of nationally representative South African data, 2008–2015. *Scientific Reports*, 10(1), 1–11.
- Tomlinson, M., Grimsrud, A. T., Stein, D. J., Williams, D. R., & Myer, L. (2009). The epidemiology of major depression in South Africa: Results from the South African stress and health study. *South African Medical Journal*, 99(5), 367–373.
- Tuthill, E. L., Maltby, A., Conte, J., Sheira, L. A., Miller, J. D., Onono, M., & Young, S. L. (2020). Persistent food insecurity, but not HIV, is associated with depressive symptoms among perinatal women in Kenya: A longitudinal perspective. *AIDS and Behavior*, 25, 847–855.
- United Nations. (2019). Over 820 million people suffering from hunger; new UN report reveals stubborn realities of 'immense' global challenge. <https://www.un.org/development/desa/en/news/sustainable/hlfp-2019-state-of-food-security-and-nutrition.html>
- Vafaei, Z., Mokhtari, H., Sadooghi, Z., Meamar, R., Chitsaz, A., & Moeini, M. (2013). Malnutrition is associated with depression in rural elderly population. *Journal of Research in Medical Sciences*, 18(suppl 1), S15.
- Whiteford, H. A., Ferrari, A. J., Degenhardt, L., Feigin, V., & Vos, T. (2015). The global burden of mental, neurological and substance use disorders: An analysis from the global burden of disease study 2010. *PLOS One*, 10(2), e0116820.
- World Health Organization. (2017). Mental health of older adults. <https://www.who.int/news-room/fact-sheets/detail/mental-health-of-older-adults>
- World Health Organization. (2020a). Depression. <https://www.who.int/health-topics/depression>
- World Health Organization. (2020b). Fact sheets: Depression. <https://www.who.int/news-room/fact-sheets/detail/depression>
- Yohannes, A. M., & Baldwin, R. C. (2008). Medical comorbidities in late-life depression. *Psychiatric Times*, 25(14), 52.
- Zimmerman, F. J., & Katon, W. (2005). Socioeconomic status, depression disparities, and financial strain: What lies behind the income-depression relationship? *Health Economics*, 14(12), 1197–1215.

**How to cite this article:** Smith, L., Il Shin, J., McDermott, D., Jacob, L., Barnett, Y., López-Sánchez, G. F., Veronese, N., Yang, L., Soysal, P., Oh, H., Grabovac, I., & Koyanagi, A.. Association between food insecurity and depression among older adults from low- and middle-income countries. *Depress Anxiety*. 2021;38:439–446. <https://doi.org/10.1002/da.23147>